



# Criterios de Autorización Previa

MEDIDUAL

# 2024

1º de enero - 31 de diciembre

Leon Health es un plan HMO que tiene contrato con Medicare.  
Inscribirse en Leon Health, Inc. depende de que se renueve el contrato

H4286\_2024PRIORAUTHCRIT002S\_C

# ABATACEPT SQ

---

## Products Affected

- ORENCIA
- ORENCIA CLICKJECT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: RHEUMATOID ARTHRITIS (RA), POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	INITIAL: RA: ONE OF THE FOLLOWING: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. PJIA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ IR. PSA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, COSENTYX, ENBREL, TREMFYA, XELJANZ, RINVOQ, SKYRIZI. RENEWAL: RA, PJIA, PSA: CONTINUES TO BENEFIT FROM THE MEDICATION.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ABEMACICLIB

---

## Products Affected

- VERZENIO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# ABIRATERONE

## Products Affected

- *abiraterone*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	METASTATIC HIGH-RISK CASTRATION-SENSITIVE PROSTATE CANCER, METASTATIC CASTRATION-RESISTANT PROSTATE CANCER: ONE OF THE FOLLOWING: 1) PREVIOUSLY RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# ABIRATERONE SUBMICRONIZED

## Products Affected

- YONSA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	METASTATIC CASTRATION-RESISTANT PROSTATE CANCER: ONE OF THE FOLLOWING: 1) PREVIOUSLY RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# ACALABRUTINIB

---

## Products Affected

- CALQUENCE
- CALQUENCE (ACALABRUTINIB MAL)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# ADAGRASIB

---

## Products Affected

- KRAZATI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# ADALIMUMAB

## Products Affected

- *adalimumab-fkjp*
- AMJEVITA(CF) SUBCUTANEOUS SYRINGE 10 MG/0.2 ML
- CYLTEZO(CF)
- CYLTEZO(CF) PEN
- CYLTEZO(CF) PEN CROHN'S-UC-HS
- CYLTEZO(CF) PEN PSORIASIS-UV
- HADLIMA
- HADLIMA PUSHTOUCH
- HADLIMA(CF)
- HADLIMA(CF) PUSHTOUCH
- HULIO(CF)
- HULIO(CF) PEN
- HUMIRA PEN
- HUMIRA PEN CROHNS-UC-HS START
- HUMIRA PEN PSOR-UEVITS-ADOL HS
- HUMIRA SUBCUTANEOUS SYRINGE KIT 40 MG/0.8 ML
- HUMIRA(CF)
- HUMIRA(CF) PEDI CROHNS STARTER
- HUMIRA(CF) PEN
- HUMIRA(CF) PEN CROHNS-UC-HS
- HUMIRA(CF) PEN PEDIATRIC UC
- HUMIRA(CF) PEN PSOR-UV-ADOL HS
- HYRIMOZ PEN CROHN'S-UC STARTER
- HYRIMOZ PEN PSORIASIS STARTER
- HYRIMOZ(CF)
- HYRIMOZ(CF) PEDI CROHN STARTER SUBCUTANEOUS SYRINGE 80 MG/0.8 ML- 40 MG/0.4 ML
- HYRIMOZ(CF) PEN
- YUSIMRY(CF) PEN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE, OR GENITAL AREA.
Age Restrictions	

PA Criteria	Criteria Details
<b>Prescriber Restrictions</b>	INITIAL: RA, POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA), ANKYLOSING SPONDYLITIS (AS): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST. UVEITIS: PRESCRIBED BY OR IN CONSULTATION WITH AN OPHTHALMOLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) - IF PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE GREATER THAN OR EQUAL TO 20 MG PER WEEK OR MAXIMALLY TOLERATED DOSE IS REQUIRED. PJIA, PSA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD. AS: TRIAL OF OR CONTRAINDICATION TO AN NSAID. PSO: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A PUVA (PHOTOTHERAPY ULTRAVIOLET LIGHT A), UVB (ULTRAVIOLET LIGHT B), TOPICAL CORTICOSTEROIDS, CALCIPOTRIEN E, ACITRETIN, METHOTREXATE, OR CYCLOSPORINE. CD, UC: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A CORTICOSTEROID (E.G., BUDESONIDE, METHYLPREDNISOLONE), AZATHIOPRINE, MERCAPTOPURINE, METHOTREXATE, OR MESALAMINE. UVEITIS: NO ISOLATED ANTERIOR UVEITIS. RENEWAL: RA, PJIA, PSA, AS, PSO, HIDRADENITIS SUPPURATIVA, UVEITIS: CONTINUES TO BENEFIT FROM THE MEDICATION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	

PA Criteria	Criteria Details
Part B Prerequisite	No

# AFATINIB

## Products Affected

- GILOTRIF

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS.
Other Criteria	METASTATIC NSCLC WITH EGFR MUTATION: NOT ON CONCURRENT THERAPY WITH AN EGFR TYROSINE KINASE-INHIBITOR.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# ALECTINIB

---

## Products Affected

- ALECENSA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# ALPELISIB

---

## Products Affected

- PIQRAY ORAL TABLET 200 MG/DAY (200 MG X 1), 250 MG/DAY (200 MG X1-50 MG X1), 300 MG/DAY (150 MG X 2)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# AMBRISENTAN

## Products Affected

- *ambrisentan*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: 1) DOCUMENTED CONFIRMATORY DIAGNOSIS BASED ON RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: A) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, B) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, AND C) PULMONARY VASCULAR RESISTANCE (PVR) OF 3 WOOD UNITS OR GREATER, AND 2) NYHA-WHO FUNCTIONAL CLASS II-IV SYMPTOMS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	PAH: INITIAL: PATIENT DOES NOT HAVE IDIOPATHIC PULMONARY FIBROSIS. RENEWAL: 1) IMPROVEMENT FROM BASELINE IN THE 6-MINUTE WALK DISTANCE, OR 2) A STABLE 6-MINUTE WALK DISTANCE WITH A STABLE/IMPROVED WHO FUNCTIONAL CLASS.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ANAKINRA

## Products Affected

- KINERET

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	CORONAVIRUS DISEASE 2019 (COVID-19) IN HOSPITALIZED ADULTS.
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: RHEUMATOID ARTHRITIS (RA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST.
<b>Coverage Duration</b>	RA: INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS. ALL OTHERS: 12 MONTHS.
<b>Other Criteria</b>	INITIAL: RA: ONE OF THE FOLLOWING: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, RINVOQ, ENBREL, XELJANZ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. RENEWAL: RA: CONTINUES TO BENEFIT FROM THE MEDICATION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# APALUTAMIDE

## Products Affected

- ERLEADA ORAL TABLET 240 MG, 60 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS.
Other Criteria	INITIAL: NON-METASTATIC CASTRATION-RESISTANT PROSTATE CANCER (NMCRPC): (1) HIGH RISK PROSTATE CANCER (I.E., RAPIDLY INCREASING PROSTATE SPECIFIC ANTIGEN [PSA] LEVELS). NMCRPC, METASTATIC CASTRATION-SENSITIVE PROSTATE CANCER (MCSPC): (1) RECEIVED A BILATERAL ORCHIECTOMY, (2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR (3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE ANALOG. RENEWAL: DIAGNOSIS OF NMCRPC OR MCSPC.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# APOMORPHINE

## Products Affected

- *apomorphine*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	PARKINSONS DISEASE (PD): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
Coverage Duration	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.
Other Criteria	PD: INITIAL: PHYSICIAN HAS OPTIMIZED DRUG THERAPY FOR PD. RENEWAL: IMPROVEMENT WITH MOTOR FLUCTUATIONS DURING OFF EPISODES WHILE ON THERAPY.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# APOMORPHINE - SL

## Products Affected

- KYNMOBI SUBLINGUAL FILM 10 MG, 10-15-20-25-30 MG, 15 MG, 20 MG, 25 MG, 30 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	PARKINSONS DISEASE (PD): INITIAL: 18 YEARS OF AGE OR OLDER.
Prescriber Restrictions	PD: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	PD: INITIAL: PHYSICIAN HAS OPTIMIZED DRUG THERAPY FOR PARKINSONS DISEASE. RENEWAL: IMPROVEMENT WITH MOTOR FLUCTUATIONS DURING OFF EPISODES WITH THE USE OF KYNMOBI.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# APREMILAST

## Products Affected

- OTEZLA
- OTEZLA STARTER

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: MILD PLAQUE PSORIASIS (PSO): ONE OF THE FOLLOWING: 1) PSORIASIS COVERING 2 PERCENT OF BODY SURFACE AREA (BSA), 2) STATIC PHYSICIAN GLOBAL ASSESSMENT (SPGA) SCORE OF 2, OR 3) PSORIASIS AREA AND SEVERITY INDEX (PASI) SCORE OF 2 TO 9. MODERATE TO SEVERE PSO: 1) PSORIASIS COVERING 3 PERCENT OR MORE OF BSA, OR 2) PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE, OR GENITAL AREA.
Age Restrictions	
Prescriber Restrictions	INITIAL: PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. BEHCETS DISEASE: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: PSA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, COSENTYX, ENBREL, TREMFYA, XELJANZ, RINVOQ, SKYRIZI. MILD PSO: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL SYSTEMIC AGENT (E.G., METHOTREXATE, ACITRETIN, CYCLOSPORINE) AND ONE CONVENTIONAL TOPICAL AGENT (E.G., PUVA, UVB, TOPICAL CORTICOSTEROIDS). MODERATE TO SEVERE PSO: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, COSENTYX, ENBREL, SKYRIZI, TREMFYA. BEHCETS DISEASE: 1) PATIENT HAS ORAL ULCERS OR A HISTORY OF RECURRENT ORAL ULCERS BASED ON CLINICAL SYMPTOMS, AND 2) TRIAL OF OR CONTRAINDICATION TO ONE OR MORE CONSERVATIVE TREATMENTS (E.G., COLCHICINE, TOPICAL CORTICOSTEROID, ORAL CORTICOSTEROID). RENEWAL: PSA, PSO, BEHCETS DISEASE: CONTINUES TO BENEFIT FROM THE MEDICATION.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ASCIMINIB

## Products Affected

- SCEMBLIX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PHILADELPHIA CHROMOSOME-POSITIVE CHRONIC MYELOID LEUKEMIA (Ph+ CML); MUTATIONAL ANALYSIS PRIOR TO INITIATION AND SCEMBLIX IS APPROPRIATE PER NCCN GUIDELINE TABLE FOR TREATMENT RECOMMENDATIONS BASED ON BCR-ABL1 MUTATION PROFILE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# ATOGEPANT

## Products Affected

- QULIPTA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	EPISODIC MIGRAINE PREVENTION: INITIAL: 1) TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING PREVENTIVE MIGRAINE TREATMENTS: DIVALPROEX SODIUM, TOPIRAMATE, PROPRANOLOL, TIMOLOL, AND 2) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION. RENEWAL: 1) REDUCTION IN MIGRAINE OR HEADACHE FREQUENCY, MIGRAINE SEVERITY, OR MIGRAINE DURATION WITH THERAPY, AND 2) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# AVAPRITINIB

---

## Products Affected

- AYVAKIT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# AVATROMBOPAG

## Products Affected

- DOPTELET (10 TAB PACK)
- DOPTELET (15 TAB PACK)
- DOPTELET (30 TAB PACK)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: CHRONIC LIVER DISEASE (CLD): PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST, GASTROENTEROLOGIST, HEPATOLOGIST, IMMUNOLOGIST, ENDOCRINOLOGIST, OR A SURGEON. CHRONIC IMMUNE THROMBOCYTOPENIA (ITP): PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR IMMUNOLOGIST.
Coverage Duration	CLD: 1 MONTH. CHRONIC ITP: INITIAL: 2 MONTHS, RENEWAL: 12 MONTHS.
Other Criteria	INITIAL: CLD: 1) PLANNED PROCEDURE 10 TO 13 DAYS AFTER INITIATION OF DOPTELET, AND 2) NOT RECEIVING OTHER THROMBOPOIETIN RECEPTOR AGONISTS (E.G., ROMIPLOSTIM, ELTROMBOPAG, ETC.). CHRONIC ITP: TRIAL OF OR CONTRAINDICATION TO CORTICOSTEROIDS OR IMMUNOGLOBULINS OR INSUFFICIENT RESPONSE TO SPLENECTOMY. RENEWAL: CHRONIC ITP: PATIENT HAD A CLINICAL RESPONSE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# AXITINIB

---

## Products Affected

- INLYTA ORAL TABLET 1 MG, 5 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# AZACITIDINE

---

## Products Affected

- ONUREG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# AZTREONAM

---

## Products Affected

- CAYSTON

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	7 YEARS OF AGE OR OLDER
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# BARICITINIB

---

## Products Affected

- OLUMIANT

PA Criteria	Criteria Details
Exclusion Criteria	CORONAVIRUS DISEASE 2019 (COVID-19) IN HOSPITALIZED ADULTS
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	RHEUMATOID ARTHRITIS (RA): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	RA: INITIAL: TRIAL OF OR CONTRAINDICATION TO ANY TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ. RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# BECAPLERMIN

## Products Affected

- REGRANEX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	DIABETIC NEUROPATHIC ULCERS: PRESCRIBED BY OR IN CONSULTATION WITH A VASCULAR SURGEON, PODIATRIST, ENDOCRINOLOGIST, PHYSICIAN PRACTICING IN A SPECIALTY WOUND CLINIC OR INFECTIOUS DISEASE SPECIALIST.
Coverage Duration	3 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# BEDAQUILINE

---

## Products Affected

- SIRTURO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	24 WEEKS
Other Criteria	PULMONARY MULTI-DRUG RESISTANT TUBERCULOSIS (MDR-TB): SIRTURO USED IN COMBINATION WITH AT LEAST 3 OTHER ANTIBIOTICS FOR THE TREATMENT OF PULMONARY MDR-TB.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# BELIMUMAB

## Products Affected

- BENLYSTA INTRAVENOUS
- BENLYSTA SUBCUTANEOUS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: SYSTEMIC LUPUS ERYTHEMATOSUS (SLE): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. LUPUS NEPHRITIS (LN): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR NEPHROLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	INITIAL: SLE: CURRENTLY TAKING CORTICOSTEROIDS, ANTIMALARIALS, NSAIDS, OR IMMUNOSUPPRESSIVE AGENTS. RENEWAL: SLE: PATIENT HAD CLINICAL IMPROVEMENT. LN: IMPROVEMENT IN RENAL RESPONSE FROM BASELINE LABORATORY VALUES (I.E., EGFR OR PROTEINURIA) AND/OR CLINICAL PARAMETERS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# BELUMOSUDIL

---

## Products Affected

- REZUROCK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# BELZUTIFAN

---

## Products Affected

- WELIREG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# BENDAMUSTINE

---

## Products Affected

- *bendamustine intravenous recon soln*
- BENDAMUSTINE INTRAVENOUS SOLUTION
- BENDEKA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# BENRALIZUMAB

---

## Products Affected

- FASENRA
- FASENRA PEN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	ASTHMA: INITIAL: BLOOD EOSINOPHIL LEVEL GREATER THAN OR EQUAL TO 150 CELLS/MCL WITHIN THE PAST 12 MONTHS.
Age Restrictions	
Prescriber Restrictions	ASTHMA: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PHYSICIAN SPECIALIZING IN ALLERGY OR PULMONARY MEDICINE.
Coverage Duration	INITIAL: 4 MONTHS, RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>ASTHMA: INITIAL: 1) CONCURRENT THERAPY WITH A MEDIUM, HIGH-DOSE, OR MAXIMALLY-TOLERATED DOSE OF AN INHALED CORTICOSTEROID (ICS) AND AT LEAST ONE OTHER MAINTENANCE MEDICATION, AND 2) ONE OF THE FOLLOWING: (A) PATIENT EXPERIENCED AT LEAST ONE ASTHMA EXACERBATION REQUIRING SYSTEMIC CORTICOSTEROID BURST LASTING 3 OR MORE DAYS WITHIN THE PAST 12 MONTHS OR AT LEAST ONE SERIOUS EXACERBATION REQUIRING HOSPITALIZATION OR ER VISIT WITHIN THE PAST 12 MONTHS, OR (B) PATIENT HAS POOR SYMPTOM CONTROL DESPITE CURRENT THERAPY AS EVIDENCED BY AT LEAST THREE OF THE FOLLOWING WITHIN THE PAST 4 WEEKS: DAYTIME ASTHMA SYMPTOMS MORE THAN TWICE/WEEK, ANY NIGHT WAKING DUE TO ASTHMA, SABA RELIEVER FOR SYMPTOMS MORE THAN TWICE/WEEK, ANY ACTIVITY LIMITATION DUE TO ASTHMA, AND 3) NOT CONCURRENTLY RECEIVING XOLAIR, DUPIXENT OR OTHER ANTI-IL5 BIOLOGICS WHEN THESE ARE USED FOR THE TREATMENT OF ASTHMA. RENEWAL: 1) NOT CONCURRENTLY RECEIVING XOLAIR, DUPIXENT OR OTHER ANTI-IL5 BIOLOGICS WHEN THESE ARE USED FOR THE TREATMENT OF ASTHMA, 2) CONTINUED USE OF ICS AND AT LEAST ONE OTHER MAINTENANCE MEDICATION, AND 3) CLINICAL RESPONSE AS EVIDENCED BY ONE OF THE FOLLOWING: (A) REDUCTION IN ASTHMA EXACERBATIONS FROM BASELINE, (B) DECREASED UTILIZATION OF RESCUE MEDICATIONS, (C) INCREASE IN PERCENT PREDICTED FEV1 FROM PRETREATMENT BASELINE, OR (D) REDUCTION IN SEVERITY OR FREQUENCY OF ASTHMA-RELATED SYMPTOMS.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# BETAINE

## Products Affected

- *betaine*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# BEXAROTENE

---

## Products Affected

- *bexarotene*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# BINIMETINIB

---

## Products Affected

- MEKTOVI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# BORTEZOMIB

## Products Affected

- *bortezomib injection recon soln 1 mg, 2.5 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# BOSENTAN

## Products Affected

- *bosentan*
- TRACLEER ORAL TABLET FOR SUSPENSION

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: 1) DOCUMENTED CONFIRMATORY DIAGNOSIS BASED ON RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: A) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, B) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, AND C) PULMONARY VASCULAR RESISTANCE (PVR) OF 3 WOOD UNITS OR GREATER, AND 2) NYHA-WHO FUNCTIONAL CLASS II-IV SYMPTOMS.
Age Restrictions	
Prescriber Restrictions	PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS.
Other Criteria	PAH: INITIAL: 1) DOES NOT HAVE ELEVATED LIVER ENZYMES (ALT, AST) MORE THAN 3 TIMES UPPER LIMIT OF NORMAL (ULN) OR INCREASE IN BILIRUBIN BY 2 OR MORE TIMES ULN, AND 2) NOT CONCURRENTLY TAKING CYCLOSPORINE A OR GLYBURIDE. RENEWAL: 1) NOT CONCURRENTLY TAKING CYCLOSPORINE A OR GLYBURIDE, 2) AGES 3 TO 17 YEARS OF AGE: A) DEMONSTRATED IMPROVEMENT IN PVR, OR B) REMAINED STABLE OR SHOWN IMPROVEMENT IN EXERCISE ABILITY, 3) AGES 18 YEARS OR OLDER: A) IMPROVEMENT FROM BASELINE IN THE 6-MINUTE WALK DISTANCE, OR B) A STABLE 6-MINUTE WALK DISTANCE WITH A STABLE/IMPROVED WHO FUNCTIONAL CLASS.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# BOSUTINIB

## Products Affected

- BOSULIF ORAL TABLET 100 MG, 400 MG, 500 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PREVIOUSLY TREATED (Ph+ CML): MUTATIONAL ANALYSIS PRIOR TO INITIATION AND BOSULIF IS APPROPRIATE PER NCCN GUIDELINE TABLE FOR TREATMENT RECOMMENDATIONS BASED ON BCR-ABL1 MUTATION PROFILE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# BRIGATINIB

## Products Affected

- ALUNBRIG ORAL TABLET 180 MG, 30 MG, 90 MG
- ALUNBRIG ORAL TABLETS,DOSE PACK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# BRONCHITOL

---

## Products Affected

- BRONCHITOL

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of Cystic fibrosis of the lung
Age Restrictions	
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist or prescribing practitioner is from a CF center accredited by the Cystic Fibrosis Foundation
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

# C1 ESTERASE INHIBITOR-HAEGARDA

## Products Affected

- HAEGARDA SUBCUTANEOUS  
RECON SOLN 2,000 UNIT, 3,000 UNIT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	HEREDITARY ANGIOEDEMA (HAE): INITIAL: DIAGNOSIS CONFIRMED BY COMPLEMENT TESTING.
Age Restrictions	
Prescriber Restrictions	HAE: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST, IMMUNOLOGIST, OR ALLERGIST.
Coverage Duration	INITIAL AND RENEWAL: 12 MONTHS.
Other Criteria	HAE: INITIAL: NOT ON CONCURRENT TREATMENT WITH ALTERNATIVE PROPHYLACTIC AGENT FOR HAE ATTACKS. RENEWAL: 1) IMPROVEMENT COMPARED TO BASELINE IN HAE ATTACKS (I.E., REDUCTIONS IN ATTACK FREQUENCY OR ATTACK SEVERITY), AND 2) NOT ON CONCURRENT TREATMENT WITH ALTERNATIVE PROPHYLACTIC AGENT FOR HAE ATTACKS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# CABOZANTINIB

## Products Affected

- COMETRIQ ORAL CAPSULE 100 MG/DAY(80 MG X1-20 MG X1), 140 MG/DAY(80 MG X1-20 MG X3), 60 MG/DAY (20 MG X 3/DAY)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# CABOZANTINIB S-MALATE - CABOMETYX

---

## Products Affected

- CABOMETYX ORAL TABLET 20 MG, 40 MG, 60 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# CANNABIDIOL

## Products Affected

- EPIDIOLEX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	DRAVET SYNDROME (DS), LENNOX-GASTAUT SYNDROME (LGS), TUBEROUS SCLEROSIS COMPLEX (TSC): PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
Coverage Duration	INITIAL: 12 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	INITIAL: LENNOX-GASTAUT SYNDROME (LGS): TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING ANTIEPILEPTIC MEDICATIONS: RUFINAMIDE, FELBAMATE, CLOBAZAM, TOPIRAMATE, LAMOTRIGINE, CLONAZEPAM. RENEWAL: DS, LGS, TSC: CONFIRMATION OF DIAGNOSIS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# CAPLACIZUMAB YHDP

## Products Affected

- CABLIVI INJECTION KIT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	ACQUIRED THROMBOTIC THROMBOCYTOPENIA PURPURA (ATTP): PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST
Coverage Duration	12 MONTHS
Other Criteria	ATTP: CABLIVI WAS PREVIOUSLY INITIATED AS PART OF THE FDA APPROVED TREATMENT REGIMEN IN COMBINATION WITH PLASMA EXCHANGE AND IMMUNOSUPPRESSIVE THERAPY WITHIN AN INPATIENT SETTING. THE PATIENT HAS NOT EXPERIENCED MORE THAN TWO RECURRENCES OF ATTP WHILE ON CABLIVI THERAPY (I.E., NEW DROP IN PLATELET COUNT REQUIRING REPEAT PLASMA EXCHANGE DURING 30 DAYS POST-PLASMA EXCHANGE THERAPY [PEX] AND UP TO 28 DAYS OF EXTENDED THERAPY).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# CAPMATINIB

---

## Products Affected

- TABRECTA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# CARGLUMIC ACID

## Products Affected

- *carglumic acid*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: ACUTE, CHRONIC HYPERAMMONEMIA (HA) DUE TO N ACETYLGLUTAMATE SYNTHASE (NAGS) DEFICIENCY: NAGS GENE MUTATION IS CONFIRMED BY BIOCHEMICAL OR GENETIC TESTING. ACUTE HA DUE TO PROPIONIC ACIDEMIA (PA): 1) CONFIRMED BY THE PRESENCE OF ELEVATED METHYLCITRIC ACID AND NORMAL METHYLMALONIC ACID, OR 2) GENETIC TESTING CONFIRMING MUTATION IN THE PCCA OR PCCB GENE. ACUTE HA DUE TO METHYLMALONIC ACIDEMIA (MMA): 1) CONFIRMED BY THE PRESENCE OF ELEVATED METHYLMALONIC ACID, METHYLCITRIC ACID, OR 2) GENETIC TESTING CONFIRMING MUTATION IN THE MMUT, MMA, MMAB OR MMADHC GENES.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	ACUTE HA DUE TO NAGS/PA/MMA: 7 DAYS. CHRONIC HA DUE TO NAGS: INITIAL: 6 MOS, RENEWAL: 12 MOS.
Other Criteria	RENEWAL: CHRONIC HA DUE TO NAGS: PATIENT HAS SHOWN CLINICAL IMPROVEMENT.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# CERITINIB

---

## Products Affected

- ZYKADIA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# CERTOLIZUMAB PEGOL

## Products Affected

- CIMZIA
- CIMZIA POWDER FOR RECONST

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE, OR GENITAL AREA. NON-RADIOGRAPHIC AXIAL SPONDYLOARTHRITIS (NR-AXSPA): 1) C-REACTIVE PROTEIN (CRP) LEVELS ABOVE THE UPPER LIMIT OF NORMAL, OR 2) SACROILIITIS ON MAGNETIC RESONANCE IMAGING (MRI).
Age Restrictions	
Prescriber Restrictions	INITIAL: RHEUMATOID ARTHRITIS (RA), ANKYLOSING SPONDYLITIS (AS), NR-AXSPA: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. CROHNS DISEASE (CD): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: RA: ONE OF THE FOLLOWING: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. PSA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, COSENTYX, ENBREL, TREMFYA, XELJANZ, RINVOQ, SKYRIZI. PSO: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, COSENTYX, ENBREL, SKYRIZI, TREMFYA. AS: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, COSENTYX, ENBREL, XELJANZ, RINVOQ. CD: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, SKYRIZI. NR-AXSPA: TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING PREFERRED AGENTS: COSENTYX, RINVOQ. PATIENTS WHO ARE PREGNANT, BREASTFEEDING, OR TRYING TO BECOME PREGNANT ARE EXCLUDED FROM STEP CRITERIA FOR ALL INDICATIONS. RENEWAL: RA, PSA, AS, PSO, NR-AXSPA: CONTINUES TO BENEFIT FROM THE MEDICATION.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# CLADRIBINE

## Products Affected

- MAVENCLAD (10 TABLET PACK)
- MAVENCLAD (4 TABLET PACK)
- MAVENCLAD (5 TABLET PACK)
- MAVENCLAD (6 TABLET PACK)
- MAVENCLAD (7 TABLET PACK)
- MAVENCLAD (8 TABLET PACK)
- MAVENCLAD (9 TABLET PACK)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL/RENEWAL: 48 WEEKS.
Other Criteria	MS: INITIAL: HAS NOT RECEIVED A TOTAL OF TWO YEARS OF MAVENCLAD TREATMENT (I.E., TWO YEARLY TREATMENT COURSES OF 2 CYCLES IN EACH). RENEWAL: 1) HAS DEMONSTRATED CLINICAL BENEFIT COMPARED TO PRE-TREATMENT BASELINE, 2) DOES NOT HAVE LYMPHOPENIA, AND 3) HAS NOT RECEIVED A TOTAL OF TWO YEARS OF MAVENCLAD TREATMENT (I.E., TWO YEARLY TREATMENT COURSES OF 2 CYCLES IN EACH).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# CLOBAZAM-SYMPAZAN

## Products Affected

- SYMPAZAN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	LENNOX-GASTAUT SYNDROME (LGS): THERAPY IS PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	LGS: 1) PATIENT IS UNABLE TO TAKE TABLETS OR SUSPENSION, AND 2) TRIAL OF OR CONTRAINDICATION TO A FORMULARY CLOBAZAM AGENT.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# COBIMETINIB

---

## Products Affected

- COTELLIC

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# CORTICOTROPIN

## Products Affected

- ACTHAR
- CORTROPHIN GEL

PA Criteria	Criteria Details
Exclusion Criteria	INITIAL: NOT APPROVED FOR DIAGNOSTIC PURPOSES.
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: ALL FDA APPROVED INDICATIONS EXCEPT INFANTILE SPASMS AND MULTIPLE SCLEROSIS (MS): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST, DERMATOLOGIST, ALLERGIST/IMMUNOLOGIST, OPHTHALMOLOGIST, PULMONOLOGIST OR NEPHROLOGIST.
Coverage Duration	INFANTILE SPASMS AND MS: 28 DAYS. OTHER FDA APPROVED INDICATIONS: INITIAL AND RENEWAL: 28 DAYS
Other Criteria	INITIAL: ALL FDA APPROVED INDICATIONS EXCEPT INFANTILE SPASMS: TRIAL OF OR CONTRAINDICATION TO INTRAVENOUS (IV) CORTICOSTEROIDS. ALL FDA APPROVED INDICATIONS EXCEPT INFANTILE SPASMS AND MS: TRIAL OF OR CONTRAINDICATION TO A STANDARD OF CARE THERAPY. RENEWAL: ALL FDA APPROVED INDICATIONS EXCEPT INFANTILE SPASMS AND MS: 1) DEMONSTRATED CLINICAL BENEFIT WHILE ON THERAPY AS INDICATED BY SYMPTOM RESOLUTION AND/OR NORMALIZATION OF LABORATORY TESTS, AND 2) CONTINUES TO POSSESS CONTRAINDICATION TO IV CORTICOSTEROIDS. PART B BEFORE PART D STEP THERAPY, APPLIES ONLY TO BENEFICIARIES IN AN MA-PD PLAN.
Indications	All FDA-approved Indications.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# CRIZANLIZUMAB-TMCA

## Products Affected

- ADAKVEO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	SICKLE CELL DISEASE (SCD): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST
Coverage Duration	INITIAL: 12 MONTHS. RENEWAL: LIFETIME
Other Criteria	SCD: INITIAL: ADULTS (18 YEARS OR OLDER): ONE OF THE FOLLOWING: (1) AT LEAST 2 SICKLE CELL CRISES IN THE PAST YEAR, (2) SICKLE-CELL ASSOCIATED SYMPTOMS WHICH ARE INTERFERING WITH ACTIVITIES OF DAILY LIVING, OR (3) HISTORY OF OR HAS RECURRENT ACUTE CHEST SYNDROME (ACS). PATIENTS BETWEEN THE AGES OF 16 TO 17 YEARS, WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA. RENEWAL FOR ALL PATIENTS: MAINTAINED OR EXPERIENCED A REDUCTION IN ACUTE COMPLICATIONS OF SICKLE CELL DISEASE. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# CRIZOTINIB

---

## Products Affected

- XALKORI ORAL CAPSULE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# CYSTEAMINE HYDROCHLORIDE

---

## Products Affected

- CYSTADROPS
- CYSTARAN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# DABRAFENIB

---

## Products Affected

- TAFINLAR ORAL CAPSULE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# DACOMITINIB

---

## Products Affected

- VIZIMPRO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	METASTATIC NSCLC: NOT ON CONCURRENT THERAPY WITH AN EGFR TYROSINE KINASE-INHIBITOR.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# DALFAMPRIDINE

## Products Affected

- *dalfampridine*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	MULTIPLE SCLEROSIS (MS): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
Coverage Duration	INITIAL: 3 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	MS: INITIAL: HAS SYMPTOMS OF A WALKING DISABILITY SUCH AS MILD TO MODERATE BILATERAL LOWER EXTREMITY WEAKNESS OR UNILATERAL WEAKNESS PLUS LOWER EXTREMITY OR TRUNCAL ATAXIA. RENEWAL: IMPROVEMENT IN WALKING ABILITY.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# DAROLUTAMIDE

## Products Affected

- NUBEQA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL AND RENEWAL: 12 MONTHS.
Other Criteria	INITIAL: NON-METASTATIC CASTRATION-RESISTANT PROSTATE CANCER (NMCRPC): PATIENT HAS HIGH RISK PROSTATE CANCER (I.E., RAPIDLY INCREASING PROSTATE SPECIFIC ANTIGEN [PSA] LEVELS) AND ONE OF THE FOLLOWING: 1) PREVIOUSLY RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG. METASTATIC HORMONE-SENSITIVE PROSTATE CANCER (MHSPC): 1) PREVIOUSLY RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG. RENEWAL: NMCRPC: NO ADDITONAL CRITERIA REQUIRED. MHSPC: 1) PREVIOUSLY RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# DASATINIB

## Products Affected

- SPRYCEL ORAL TABLET 100 MG, 140 MG, 20 MG, 50 MG, 70 MG, 80 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PREVIOUSLY TREATED Ph+ CML: MUTATIONAL ANALYSIS PRIOR TO INITIATION AND SPRYCEL IS APPROPRIATE PER NCCN GUIDELINE TABLE FOR TREATMENT RECOMMENDATIONS BASED ON BCR-ABL1 MUTATION PROFILE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# DAYBUE

---

## Products Affected

- DAYBUE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of Rett syndrome
Age Restrictions	2 years of age and older
Prescriber Restrictions	
Coverage Duration	6 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

# DECITABINE/CEDAZURIDINE

---

## Products Affected

- INQOVI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# DEFERASIROX

## Products Affected

- *deferasirox*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: CHRONIC IRON OVERLOAD DUE TO BLOOD TRANSFUSIONS: SERUM FERRITIN LEVEL CONSISTENTLY ABOVE 1000 MCG/L (AT LEAST TWO LAB VALUES IN THE PREVIOUS THREE MONTHS). NON-TRANSFUSION DEPENDENT THALASSEMIA (NTDT): 1) SERUM FERRITIN LEVEL CONSISTENTLY ABOVE 300 MCG/L (AT LEAST TWO LAB VALUES IN THE PREVIOUS THREE MONTHS) AND 2) LIVER IRON CONCENTRATION (LIC) OF 5 MG FE/G DRY WEIGHT OR GREATER. RENEWAL: CHRONIC IRON OVERLOAD DUE TO BLOOD TRANSFUSIONS: SERUM FERRITIN LEVEL CONSISTENTLY ABOVE 500 MCG/L (AT LEAST TWO LAB VALUES IN THE PREVIOUS THREE MONTHS). NTDT: 1) SERUM FERRITIN LEVEL CONSISTENTLY ABOVE 300 MCG/L (AT LEAST TWO LAB VALUES IN THE PREVIOUS THREE MONTHS) OR 2) LIC OF 3 MG FE/G DRY WEIGHT OR GREATER.
Age Restrictions	
Prescriber Restrictions	INITIAL: CHRONIC IRON OVERLOAD: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR HEMATOLOGIST/ONCOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	INITIAL FOR ALL INDICATIONS: FORMULARY VERSION OF DEFERASIROX SPRINKLE: TRIAL OF OR CONTRAINDICATION TO GENERIC DEFERASIROX TABLET OR TABLET FOR ORAL SUSPENSION.
Indications	All FDA-approved Indications.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# DEFERIPRONE

## Products Affected

- *deferiprone*
- FERRIPROX (2 TIMES A DAY)
- FERRIPROX ORAL SOLUTION

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	TRANSFUSIONAL IRON OVERLOAD: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR HEMATOLOGIST/ONCOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	INITIAL: TRANSFUSIONAL IRON OVERLOAD DUE TO THALASSEMIA SYNDROMES: 1) TRIAL OF, CONTRAINDICATION, INTOLERABLE TOXICITIES, OR CLINICALLY SIGNIFICANT ADVERSE EFFECTS TO ONE OF THE FOLLOWING: FORMULARY VERSION OF DEFERASIROX OR DEFEROXAMINE, OR 2) CURRENT CHELATION THERAPY (I.E., FORMULARY VERSION OF DEFERASIROX OR DEFEROXAMINE) IS INADEQUATE. TRANSFUSIONAL IRON OVERLOAD DUE TO SICKLE CELL DISEASE OR OTHER ANEMIAS: TRIAL OF OR CONTRAINDICATION TO A FORMULARY VERSION OF DEFERASIROX OR DEFEROXAMINE. RENEWAL (ALL INDICATIONS): SERUM FERRITIN LEVELS CONSISTENTLY ABOVE 500MCG/L (AT LEAST TWO LAB VALUES IN THE PREVIOUS THREE MONTHS).
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

# DEFEROXAMINE

## Products Affected

- *deferoxamine injection recon soln 500 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	CHRONIC IRON OVERLOAD: 3 YEARS OR OLDER
Prescriber Restrictions	CHRONIC IRON OVERLOAD: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR HEMATOLOGIST/ONCOLOGIST
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	CHRONIC IRON OVERLOAD: INITIAL: SERUM FERRITIN LEVEL CONSISTENTLY ABOVE 1000MCG/L (AT LEAST TWO LAB VALUES IN THE PREVIOUS THREE MONTHS). RENEWAL: SERUM FERRITIN LEVELS MUST BE CONSISTENTLY ABOVE 500MCG/L (AT LEAST TWO LAB VALUES IN THE PREVIOUS THREE MONTHS). THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. THIS DRUG MAY BE EITHER BUNDLED WITH AND COVERED UNDER END STAGE RENAL DISEASE DIALYSIS RELATED SERVICES OR COVERED UNDER MEDICARE D DEPENDING UPON THE CIRCUMSTANCES. INFORMATION MAY NEED TO BE SUBMITTED DESCRIBING THE USE AND SETTING OF THE DRUG TO MAKE THE DETERMINATION.
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

# DENOSUMAB-XGEVA

---

## Products Affected

- XGEVA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# DEUTETRABENAZINE

## Products Affected

- AUSTEDO ORAL TABLET 12 MG, 6 MG, 9 MG
- AUSTEDO XR ORAL TABLET EXTENDED RELEASE 24 HR 12 MG, 24 MG, 6 MG
- AUSTEDO XR TITRATION KT(WK1-4)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	HUNTINGTON DISEASE: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST OR MOVEMENT DISORDER SPECIALIST. TARDIVE DYSKINESIA: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST, PSYCHIATRIST, OR MOVEMENT DISORDER SPECIALIST.
Coverage Duration	12 MONTHS
Other Criteria	TARDIVE DYSKINESIA: PATIENT HAS A HISTORY OF USING AGENTS THAT CAUSE TARDIVE DYSKINESIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# DICLOFENAC GEL

---

## Products Affected

- *diclofenac sodium topical gel 3 %*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# DICLOFENAC TOPICAL SOLUTION

## Products Affected

- *diclofenac sodium topical solution in metered-dose pump*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 MONTHS
Other Criteria	OSTEOARTHRITIS OF THE KNEE: TRIAL OF OR CONTRAINDICATION TO A FORMULARY VERSION OF DICLOFENAC SODIUM 1% TOPICAL GEL AND A FORMULARY VERSION OF DICLOFENAC SODIUM 1.5% TOPICAL DROPS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# DIMETHYL FUMARATE

## Products Affected

- *dimethyl fumarate oral capsule, delayed release(dr/ec) 120 mg, 120 mg (14)- 240 mg (46), 240 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# DIROXIMEL FUMARATE

---

## Products Affected

- VUMERITY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# DRONABINOL

## Products Affected

- *dronabinol*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 MONTHS
Other Criteria	NAUSEA AND VOMITING ASSOCIATED WITH CANCER CHEMOTHERAPY: TRIAL OF OR CONTRAINDICATION TO ONE ANTIEMETIC THERAPY. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D FOR THE INDICATION OF NAUSEA AND VOMITING ASSOCIATED WITH CANCER CHEMOTHERAPY.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# DROXIDOPA

## Products Affected

- *droxidopa*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	NEUROGENIC ORTHOSTATIC HYPOTENSION (NOH): INITIAL: 1) BASELINE BLOOD PRESSURE READINGS WHILE THE PATIENT IS SITTING AND ALSO WITHIN 3 MINUTES OF STANDING FROM A SUPINE POSITION. 2) A DECREASE OF AT LEAST 20 MMHG IN SYSTOLIC BLOOD PRESSURE OR 10 MMHG DIASTOLIC BLOOD PRESSURE WITHIN THREE MINUTES AFTER STANDING FROM A SITTING POSITION.
Age Restrictions	
Prescriber Restrictions	NOH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST OR CARDIOLOGIST.
Coverage Duration	INITIAL: 3 MONTHS RENEWAL: 12 MONTHS
Other Criteria	NOH: RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# DUPILUMAB

## Products Affected

- DUPIXENT PEN
- DUPIXENT SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: EOSINOPHILIC ASTHMA: BLOOD EOSINOPHIL LEVEL GREATER THAN OR EQUAL TO 150 CELLS/MCL WITHIN THE PAST 12 MONTHS. EOSINOPHILIC ESOPHAGITIS (EOE): DIAGNOSIS CONFIRMED BY ESOPHAGOGASTRODUODENOSCOPY (EGD) WITH BIOPSY.
Age Restrictions	
Prescriber Restrictions	INITIAL: AD, PN: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST, ALLERGIST OR IMMUNOLOGIST. ASTHMA: PRESCRIBED BY OR IN CONSULTATION WITH A PHYSICIAN SPECIALIZING IN ALLERGY OR PULMONARY MEDICINE. CRSWNP: PRESCRIBED BY OR IN CONSULTATION WITH AN OTOLARYNGOLOGIST, ALLERGIST OR IMMUNOLOGIST. EOE: PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST, ALLERGIST, OR IMMUNOLOGIST.
Coverage Duration	INITIAL: AD, CRSWNP, EOE, PN: 6 MOS, ASTHMA: 4 MOS. RENEWAL: ALL INDICATIONS: 12 MOS.

PA Criteria	Criteria Details
Other Criteria	<p>INITIAL: AD: 1) AD COVERING AT LEAST 10 PERCENT OF BODY SURFACE AREA OR AD AFFECTING THE FACE, HEAD, NECK, HANDS, FEET, GROIN, OR INTERTRIGINOUS AREAS, 2) INTRACTABLE PRURITUS OR CRACKING/OOZING/BLEEDING OF AFFECTED SKIN, 3) TRIAL OF OR CONTRAINDICATION TO ONE TOPICAL (CORTICOSTEROID, CALCINEURIN INHIBITOR, PDE4 INHIBITOR, OR JAK INHIBITOR), AND 4) NO CONCURRENT USE WITH OTHER SYSTEMIC BIOLOGIC/JAK INHIBITOR FOR AD. ASTHMA: 1) CONCURRENT THERAPY WITH A MEDIUM, HIGH-DOSE OR MAXIMALLY-TOLERATED DOSE OF AN INHALED CORTICOSTEROID (ICS) AND ONE OTHER MAINTENANCE MEDICATION, 2) ONE ASTHMA EXACERBATION REQUIRING SYSTEMIC CORTICOSTEROID BURST LASTING 3 OR MORE DAYS WITHIN THE PAST 12 MONTHS, OR ONE SERIOUS EXACERBATION REQUIRING HOSPITALIZATION OR ER VISIT WITHIN THE PAST 12 MONTHS, OR POOR SYMPTOM CONTROL DESPITE CURRENT THERAPY AS EVIDENCED BY AT LEAST THREE OF THE FOLLOWING WITHIN THE PAST 4 WEEKS: DAYTIME ASTHMA SYMPTOMS MORE THAN TWICE/WEEK, ANY NIGHT WAKING DUE TO ASTHMA, SABA RELIEVER FOR SYMPTOMS MORE THAN TWICE/WEEK, ANY ACTIVITY LIMITATION DUE TO ASTHMA, AND 3) NO CONCURRENT USE WITH XOLAIR OR OTHER ANTI-IL5 BIOLOGICS WHEN USED FOR ASTHMA. CRSWNP: 1) EVIDENCE OF NASAL POLYPS BY DIRECT EXAMINATION, ENDOSCOPY OR SINUS CT SCAN, 2) INADEQUATELY CONTROLLED DISEASE AS DETERMINED BY USE OF SYSTEMIC STEROIDS IN THE PAST 2 YEARS OR ENDOSCOPIC SINUS SURGERY, AND 3) A 90 DAY TRIAL OF ONE TOPICAL NASAL CORTICOSTEROID. PN: 1) CHRONIC PRURITIS (ITCH MORE</p>



PA Criteria	Criteria Details
	<p>THAN 6 WEEKS), MULTIPLE PRURIGINOUS LESIONS, AND HISTORY OR SIGN OF A PROLONGED SCRATCHING BEHAVIOR, 2) TRIAL OF OR CONTRAINDICATION TO ONE TOPICAL (CORTICOSTEROID OR CALCIPOTRIOL).</p> <p>RENEWAL: AD: 1) IMPROVEMENT WHILE ON THERAPY, AND 2) NO CONCURRENT USE WITH OTHER SYSTEMIC BIOLOGIC/JAK INHIBITOR FOR AD. CRSWNP, EOE: IMPROVEMENT WHILE ON THERAPY. ASTHMA: 1) NO CONCURRENT USE WITH XOLAIR, OR OTHER ANTI-IL5 BIOLOGICS FOR ASTHMA, 2) CONTINUED USE OF ICS AND ONE OTHER MAINTENANCE MEDICATION, AND 3) CLINICAL RESPONSE AS EVIDENCED BY: (A) REDUCTION IN ASTHMA EXACERBATIONS FROM BASELINE, (B) DECREASED UTILIZATION OF RESCUE MEDICATIONS, (C) INCREASE IN PERCENT PREDICTED FEV1 FROM PRETREATMENT BASELINE, OR (D) REDUCTION IN SEVERITY OR FREQUENCY OF ASTHMA-RELATED SYMPTOMS. PN: IMPROVEMENT OR REDUCTION OF PRURITIS OR PRURIGINOUS LESIONS.</p>
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# DUVELISIB

---

## Products Affected

- COPIKTRA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# EDARAVONE

## Products Affected

- RADICAVA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	AMYOTROPHIC LATERAL SCLEROSIS (ALS): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST OR ALS SPECIALIST AT AN ALS SPECIALTY CENTER OR CARE CLINIC.
Coverage Duration	ALS: INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.
Other Criteria	ALS: INITIAL: 1) DURATION OF DISEASE (FROM ONSET OF SYMPTOMS) IS LESS THAN OR EQUAL TO 2 YEARS, 2) NORMAL RESPIRATORY FUNCTION, 3) HAS MILD TO MODERATE ALS WITH A SCORE OF 2 OR HIGHER IN ALL OF THE FOLLOWING 12 ITEMS OF THE AMYOTROPHIC LATERAL SCLEROSIS FUNCTIONAL RATING SCALE REVISED (ALSFRS-R): SPEECH, SALIVATION, SWALLOWING, HANDWRITING, CUTTING FOOD, DRESSING AND HYGIENE, TURNING IN BED, WALKING, CLIMBING STAIRS, DYSPNEA, ORTHOPNEA, RESPIRATORY INSUFFICIENCY, AND 4) TRIAL OF RILUZOLE TABLET OR CURRENTLY TAKING RILUZOLE TABLET. RENEWAL: 1) DOES NOT REQUIRE INVASIVE VENTILATION, AND 2) HAS IMPROVED BASELINE FUNCTIONAL ABILITY OR HAS MAINTAINED A SCORE OF 2 OR HIGHER IN ALL 12 ITEMS OF THE ALSFRS-R.
Indications	All FDA-approved Indications.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ELACESTRANT

---

## Products Affected

- ORSERDU ORAL TABLET 345 MG, 86 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# ELAGOLIX SODIUM

## Products Affected

- ORILISSA ORAL TABLET 150 MG, 200 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	MODERATE TO SEVERE PAIN ASSOCIATED WITH ENDOMETRIOSIS: INITIAL: DIAGNOSIS IS CONFIRMED VIA SURGICAL OR DIRECT VISUALIZATION (E.G., PELVIC ULTRASOUND) OR HISTOPATHOLOGICAL CONFIRMATION (E.G., LAPAROSCOPY OR LAPAROTOMY) IN THE LAST 10 YEARS.
<b>Age Restrictions</b>	MODERATE TO SEVERE PAIN ASSOCIATED WITH ENDOMETRIOSIS: INITIAL: 18 YEARS OF AGE OR OLDER.
<b>Prescriber Restrictions</b>	MODERATE TO SEVERE PAIN ASSOCIATED WITH ENDOMETRIOSIS: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN OBSTETRICIAN/GYNECOLOGIST.
<b>Coverage Duration</b>	MODERATE TO SEVERE PAIN ASSOCIATED WITH ENDOMETRIOSIS: INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	MODERATE TO SEVERE PAIN ASSOCIATED WITH ENDOMETRIOSIS: INITIAL: 1) NO CONCURRENT USE WITH ANOTHER GNRH-MODULATING AGENT, AND 2) TRIAL OF OR CONTRAINDICATION TO A NSAID AND PROGESTIN-CONTAINING PREPARATION. RENEWAL: 1) IMPROVEMENT IN PAIN ASSOCIATED WITH ENDOMETRIOSIS WHILE ON THERAPY, AND 2) NO CONCURRENT USE WITH ANOTHER GNRH-MODULATING AGENT.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	

PA Criteria	Criteria Details
Part B Prerequisite	No

# ELAPEGADEMASE-LVLR

## Products Affected

- REVCovi

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	ADENOSINE DEAMINASE SEVERE COMBINED IMMUNE DEFICIENCY (ADA-SCID): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH IMMUNOLOGIST, HEMATOLOGIST/ONCOLOGIST, OR PHYSICIAN SPECIALIZING IN INHERITED METABOLIC DISORDERS
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	ADA-SCID: INITIAL: ADA-SCID AS MANIFESTED BY ONE OF THE FOLLOWING: (1) CONFIRMATORY GENETIC TEST OR (2) SUGGESTIVE LABORATORY FINDINGS (E.G., ELEVATED DEOXYADENOSINE NUCLEOTIDE [DAXP] LEVELS, LYMPHOPENIA) AND HALLMARK SIGNS/SYMPTOMS (E.G., RECURRENT INFECTIONS, FAILURE TO THRIVE, PERSISTENT DIARRHEA). RENEWAL: 1) IMPROVEMENT OR MAINTENANCE OF IMMUNE FUNCTION FROM BASELINE, AND 2) HAS NOT RECEIVED SUCCESSFUL HCT OR GENE THERAPY.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# ELEXACFTOR-TEZACFTOR-IVACFTOR

## Products Affected

- TRIKAFTA ORAL TABLETS, SEQUENTIAL

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	CYSTIC FIBROSIS (CF): INITIAL: CONFIRMED MUTATION IN CFTR GENE ACCEPTABLE FOR THE TREATMENT OF CYSTIC FIBROSIS.
Age Restrictions	
Prescriber Restrictions	CF: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST OR CYSTIC FIBROSIS EXPERT.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: LIFETIME.
Other Criteria	CF: RENEWAL: 1) MAINTAINED, IMPROVED, OR DEMONSTRATED LESS THAN EXPECTED DECLINE IN FEV1 OR BODY MASS INDEX (BMI), OR 2) REDUCTION IN NUMBER OF PULMONARY EXACERBATIONS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# ELIGLUSTAT

---

## Products Affected

- CERDELGA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# ELTROMBOPAG

## Products Affected

- PROMACTA ORAL POWDER IN PACKET 12.5 MG, 25 MG
- PROMACTA ORAL TABLET 12.5 MG, 25 MG, 50 MG, 75 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: ITP: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR IMMUNOLOGIST.
Coverage Duration	ITP: INITIAL: 2 MO, RENEWAL: 12 MO. HEPATITIS C, SEVERE APLASTIC ANEMIA: 12 MO.
Other Criteria	INITIAL: PERSISTENT OR CHRONIC IMMUNE THROMBOCYTOPENIA PURPURA (ITP): TRIAL OF OR CONTRAINDICATION TO CORTICOSTEROIDS, IMMUNOGLOBULINS, OR AN INSUFFICIENT RESPONSE TO SPLENECTOMY. ALL INDICATIONS: APPROVAL FOR PROMACTA ORAL SUSPENSION PACKETS REQUIRES A TRIAL OF PROMACTA TABLETS OR PATIENT IS UNABLE TO TAKE TABLET FORMULATION. RENEWAL: ITP: PATIENT HAS SHOWN A CLINICAL RESPONSE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# ENASIDENIB

---

## Products Affected

- IDHIFA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# ENCORAFENIB

---

## Products Affected

- BRAFTOVI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# ENTRECTINIB

---

## Products Affected

- ROZLYTREK ORAL CAPSULE 100 MG, 200 MG
- ROZLYTREK ORAL PELLETS IN PACKET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# ENZALUTAMIDE

## Products Affected

- XTANDI ORAL CAPSULE
- XTANDI ORAL TABLET 40 MG, 80 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	INITIAL: CASTRATION-RESISTANT PROSTATE CANCER (CRPC) THAT IS NOT METASTATIC: PATIENT HAS HIGH RISK PROSTATE CANCER (I.E. RAPIDLY INCREASING PROSTATE SPECIFIC ANTIGEN [PSA] LEVELS). CRPC (INCLUDES NON-METASTATIC AND METASTATIC) OR METASTATIC CASTRATION-SENSITIVE PROSTATE CANCER (MCSPC): 1) PREVIOUSLY RECEIVED A BILATERAL ORCHIECTOMY, OR 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG. RENEWAL: DIAGNOSIS OF CRPC (INCLUDES NON-METASTATIC AND METASTATIC) OR MCSPC.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# EPOETIN ALFA-EPBX

## Products Affected

- RETACRIT INJECTION SOLUTION  
10,000 UNIT/ML, 2,000 UNIT/ML,  
20,000 UNIT/2 ML, 20,000 UNIT/ML,  
3,000 UNIT/ML, 4,000 UNIT/ML, 40,000  
UNIT/ML

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: CHRONIC KIDNEY DISEASE (CKD), ANEMIA RELATED TO ZIDOVUDINE, OR CANCER CHEMOTHERAPY: HEMOGLOBIN LEVEL OF LESS THAN 10G/DL. ELECTIVE, NON-CARDIAC, NON-VASCULAR SURGERY: HEMOGLOBIN LEVEL LESS THAN 13G/DL. RENEWAL: CKD: 1) HEMOGLOBIN LEVEL IS LESS THAN 10G/DL, OR 2) HEMOGLOBIN LEVEL HAS REACHED 10G/DL AND DOSE REDUCTION/INTERRUPTION IS REQUIRED TO REDUCE THE NEED FOR BLOOD TRANSFUSIONS. ANEMIA RELATED TO ZIDOVUDINE: HEMOGLOBIN LEVEL BETWEEN 10G/DL AND 12G/DL. CANCER CHEMOTHERAPY: 1) HEMOGLOBIN LEVEL OF LESS THAN 10 G/DL, OR 2) THE HEMOGLOBIN LEVEL DOES NOT EXCEED A LEVEL NEEDED TO AVOID RBC TRANSFUSION.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	ANEMIA FROM CHEMO/CKD WITHOUT DIALYSIS/ZIDOVUDINE: INITIAL/RENEWAL: 12 MONTHS. SURGERY: 1 MONTH.
<b>Other Criteria</b>	RENEWAL: CKD: NOT RECEIVING DIALYSIS TREATMENT. THIS DRUG MAY BE EITHER BUNDLED WITH AND COVERED UNDER END STAGE RENAL DISEASE DIALYSIS RELATED SERVICES OR COVERED UNDER MEDICARE D DEPENDING UPON THE CIRCUMSTANCES. INFORMATION MAY NEED TO BE SUBMITTED DESCRIBING THE USE AND SETTING OF THE DRUG TO MAKE THE DETERMINATION.



<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# EPOPROSTENOL IV

## Products Affected

- *epoprostenol*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: (1) DOCUMENTED CONFIRMATORY PAH DIAGNOSIS BASED ON RIGHT HEART CATHETERIZATION: A) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, B) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) 15 MMHG OR LESS, AND C) PULMONARY VASCULAR RESISTANCE (PVR) 3 WOOD UNITS OR GREATER, AND (2) NYHA-WHO FUNCTIONAL CLASS III-IV SYMPTOMS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
<b>Coverage Duration</b>	INITIAL AND RENEWAL: 12 MONTHS
<b>Other Criteria</b>	PAH: RENEWAL: IMPROVEMENT FROM BASELINE IN THE 6-MINUTE WALK DISTANCE OR A STABLE 6-MINUTE WALK DISTANCE WITH A STABLE/IMPROVED WHO FUNCTIONAL CLASS. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ERDAFITINIB

---

## Products Affected

- BALVERSA ORAL TABLET 3 MG, 4 MG, 5 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# ERLOTINIB

## Products Affected

- *erlotinib oral tablet 100 mg, 150 mg, 25 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	METASTATIC NSCLC WITH EGFR MUTATION: NOT ON CONCURRENT THERAPY WITH AN EGFR TYROSINE KINASE-INHIBITOR.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# ESKETAMINE

## Products Affected

- SPRAVATO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: TREATMENT-RESISTANT DEPRESSION (TRD), MAJOR DEPRESSIVE DISORDER (MDD): PRESCRIBED BY OR IN CONSULTATION WITH A PSYCHIATRIST.
Coverage Duration	INITIAL: TRD: 3 MONTHS. MDD: 4 WEEKS. RENEWAL: TRD, MDD: 12 MONTHS.
Other Criteria	INITIAL: TRD: 1) NON-PSYCHOTIC, UNIPOLAR DEPRESSION, 2) NO ACTIVE SUBSTANCE ABUSE, AND 3) ADEQUATE TRIAL (AT LEAST 4 WEEKS) OF AT LEAST TWO ANTIDEPRESSANT AGENTS FROM DIFFERENT CLASSES THAT ARE INDICATED FOR DEPRESSION. MDD: 1) NON-PSYCHOTIC, UNIPOLAR DEPRESSION, AND 2) NO ACTIVE SUBSTANCE ABUSE. RENEWAL: TRD, MDD: DEMONSTRATED CLINICAL BENEFIT (IMPROVEMENT IN DEPRESSION) COMPARED TO BASELINE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# ETANERCEPT

## Products Affected

- ENBREL
- ENBREL MINI
- ENBREL SURECLICK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE OR GENITAL AREA.
Age Restrictions	INITIAL: RHEUMATOID ARTHRITIS (RA), ANKYLOSING SPONDYLITIS (AS), PSORIATIC ARTHRITIS (PSA): 18 YEARS OR OLDER.
Prescriber Restrictions	INITIAL: RA, POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA), AS: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSA: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Other Criteria</b>	INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) - IF PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE GREATER THAN OR EQUAL TO 20 MG PER WEEK OR MAXIMALLY TOLERATED DOSE: IS REQUIRED. PJIA, PSA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD. AS: TRIAL OF OR CONTRAINDICATION TO AN NSAID. PSO: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A PUVA (PHOTOTHERAPY ULTRAVIOLET LIGHT A), UVB (ULTRAVIOLET LIGHT B), TOPICAL CORTICOSTEROIDS, CALCIPOTRIENE, ACITRETIN, METHOTREXATE, OR CYCLOSPORINE. RENEWAL: RA, PJIA, PSA, AS, PSO: CONTINUES TO BENEFIT FROM THE MEDICATION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# EVEROLIMUS

## Products Affected

- *everolimus (antineoplastic) oral tablet 10 mg, 2.5 mg, 5 mg, 7.5 mg*
- *everolimus (antineoplastic) oral tablet for suspension*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# FEDRATINIB

## Products Affected

- INREBIC

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	MYELOFIBROSIS: INITIAL: TRIAL OF OR CONTRAINDICATION TO JAKAFI (RUXOLITINIB). RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# FENFLURAMINE

## Products Affected

- FINTEPLA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: DRAVET SYNDROME, LENNOX-GASTAUT SYNDROME (LGS): PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
Coverage Duration	DRAVET SYNDROME: INITIAL/RENEWAL: 12 MONTHS. LGS: 12 MONTHS.
Other Criteria	INITIAL: LGS: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING ANTIEPILEPTIC MEDICATIONS: RUFINAMIDE, FELBAMATE, CLOBAZAM, TOPIRAMATE, LAMOTRIGINE, CLONAZEPAM. RENEWAL: DRAVET SYNDROME: PATIENT HAS SHOWN CONTINUED CLINICAL BENEFIT (E.G. REDUCTION OF SEIZURES, REDUCED LENGTH OF SEIZURES, SEIZURE CONTROL MAINTAINED).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# FENTANYL CITRATE

## Products Affected

- *fentanyl citrate buccal lozenge on a handle*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	CANCER RELATED PAIN: 1) CURRENTLY ON A MAINTENANCE DOSE OF CONTROLLED-RELEASE OPIOID PAIN MEDICATION, AND 2) TRIAL OF OR CONTRAINDICATION TO AT LEAST ONE IMMEDIATE-RELEASE ORAL OPIOID PAIN AGENT OR PATIENT HAS DIFFICULTY SWALLOWING TABLETS/CAPSULES. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# FILGRASTIM-AAFI

---

## Products Affected

- NIVESTYM

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# FILGRASTIM-AYOW

## Products Affected

- RELEUKO SUBCUTANEOUS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	TRIAL OF OR CONTRAINDICATION TO THE PREFERRED AGENT NIVESTYM, WHERE INDICATIONS ALIGN.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# FILGRASTIM-SNDZ

---

## Products Affected

- ZARXIO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	TRIAL OF OR CONTRAINDICATION TO THE PREFERRED AGENT: NIVESTYM.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# FINASTERIDE/TADALAFIL

---

## Products Affected

- ENTADFI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	BENIGN PROSTATIC HYPERPLASIA (BPH): 18 YEARS OF AGE OR OLDER
Prescriber Restrictions	
Coverage Duration	26 WEEKS
Other Criteria	BPH: 1) TRIAL OF OR CONTRAINDICATION TO ONE 5-ALPHA-REDUCTASE INHIBITOR, AND 2) TRIAL OF OR CONTRAINDICATION TO TADALAFIL 2.5MG OR TADALAFIL 5MG.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# FINERENONE

---

## Products Affected

- KERENDIA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# FINGOLIMOD

---

## Products Affected

- *fingolimod*
- GILENYA ORAL CAPSULE 0.25 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# FINGOLIMOD LAURYL SULFATE

---

## Products Affected

- TASCENSO ODT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	MULTIPLE SCLEROSIS (MS): (1) UNABLE TO SWALLOW FINGOLIMOD CAPSULES, AND (2) TRIAL OF OR CONTRAINDICATION TO FINGOLIMOD CAPSULES.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# FIRMAGON

## Products Affected

- FIRMAGON KIT W DILUENT  
SYRINGE SUBCUTANEOUS RECON  
SOLN 120 MG, 80 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of advanced prostate cancer
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

# FOSTAMATINIB

## Products Affected

- TAVALISSE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	CHRONIC IMMUNE THROMBOCYTOPENIA (ITP): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR IMMUNOLOGIST.
Coverage Duration	INITIAL: 3 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	ITP: RENEWAL: PATIENT HAS SHOWN A CLINICAL RESPONSE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# FREMANEZUMAB-VFRM

## Products Affected

- AJOVY AUTOINJECTOR
- AJOVY SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.
Other Criteria	MIGRAINE PREVENTION: INITIAL: 1) TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING PREVENTIVE MIGRAINE TREATMENTS: DIVALPROEX SODIUM, TOPIRAMATE, PROPRANOLOL, TIMOLOL, AND 2) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION. RENEWAL: 1) REDUCTION IN MIGRAINE OR HEADACHE FREQUENCY, MIGRAINE SEVERITY, OR MIGRAINE DURATION WITH THERAPY, AND 2) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# FRUZAQLA

---

## Products Affected

- FRUZAQLA ORAL CAPSULE 1 MG, 5 MG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of metastatic colorectal cancer (mCRC) and all of the following: A.) patient has been previously treated with fluoropyrimidine, oxaliplatin, irinotecan-based chemotherapy, B.) an anti-VEGF therapy, and C.) if RAS wild-type and medically appropriate, patient has also been previously treated with anti-EGFR therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	3 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

# FUTIBATINIB

## Products Affected

- LYTGOBI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INTRAHEPATIC CHOLANGIOCARCINOMA (ICCA): COMPLETE A COMPREHENSIVE OPHTHALMOLOGICAL EXAMINATION, INCLUDING OPTICAL COHERENCE TOMOGRAPHY (OCT), PRIOR TO THE INITIATION OF THERAPY AND AT THE RECOMMENDED SCHEDULED INTERVALS.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# GALCANEZUMAB-GNLM

## Products Affected

- EMGALITY PEN 3)
- EMGALITY SYRINGE
- SUBCUTANEOUS SYRINGE 120 MG/ML, 300 MG/3 ML (100 MG/ML X

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL: MIGRAINE PREVENTION: 6 MOS. EPISODIC CLUSTER HEADACHE: 3 MOS. RENEWAL (ALL): 12 MOS.
Other Criteria	INITIAL: MIGRAINE PREVENTION: 1) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION, AND 2) TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING PREVENTIVE MIGRAINE TREATMENTS: DIVALPROEX SODIUM, TOPIRAMATE, PROPRANOLOL, TIMOLOL. RENEWAL: MIGRAINE PREVENTION: 1) REDUCTION IN MIGRAINE OR HEADACHE FREQUENCY, MIGRAINE SEVERITY, OR MIGRAINE DURATION WITH THERAPY, AND 2) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION. EPISODIC CLUSTER HEADACHE: IMPROVEMENT IN EPISODIC CLUSTER HEADACHE FREQUENCY AS COMPARED TO BASELINE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# GANAXOLONE

---

## Products Affected

- ZTALMY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# GEFITINIB

## Products Affected

- *gefitinib*
- IRESSA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	METASTATIC NSCLC WITH EGFR MUTATION: NOT ON CONCURRENT THERAPY WITH AN EGFR TYROSINE KINASE-INHIBITOR.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# GILTERITINIB

---

## Products Affected

- XOSPATA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# GLASDEGIB

---

## Products Affected

- DAURISMO ORAL TABLET 100 MG, 25 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# GLATIRAMER

## Products Affected

- COPAXONE SUBCUTANEOUS SYRINGE 20 MG/ML, 40 MG/ML
- *glatiramer subcutaneous syringe 20 mg/ml, 40 mg/ml*
- *glatopa subcutaneous syringe 20 mg/ml, 40 mg/ml*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# GLECAPREVIR/PIBRENTASVIR

## Products Affected

- MAVYRET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	HCV RNA LEVEL WITHIN PAST 6 MONTHS
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE.
Other Criteria	1) CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE, 2) TRIAL OF A PREFERRED FORMULARY ALTERNATIVE INCLUDING HARVONI OR EPCLUSA WHEN THESE AGENTS ARE CONSIDERED ACCEPTABLE FOR TREATMENT OF THE SPECIFIC GENOTYPE PER AASLD/IDSA GUIDANCE, 3) NOT CONCURRENTLY TAKING ANY OF THE FOLLOWING MEDICATIONS: CARBAMAZEPINE, RIFAMPIN, ETHINYL ESTRADIOL-CONTAINING MEDICATION, ATAZANAVIR, DARUNAVIR, LOPINAVIR, RITONAVIR, EFAVIRENZ, ATORVASTATIN, LOVASTATIN, SIMVASTATIN, ROSUVASTATIN AT DOSES GREATER THAN 10MG, CYCLOSPORINE AT DOSES GREATER THAN 100MG PER DAY, EPCLUSA, HARVONI, VOSEVI, OR ZEPATIER, 4) PATIENT MUST NOT HAVE PRIOR FAILURE OF A DAA REGIMEN WITH NS5A INHIBITOR AND HCV PROTEASE INHIBITOR, AND 5) DOES NOT HAVE MODERATE OR SEVERE HEPATIC IMPAIRMENT (CHILD PUGH B OR C).
Indications	All FDA-approved Indications.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# GLP1 AGONISTS

## Products Affected

- OZEMPIC SUBCUTANEOUS PEN INJECTOR 1 MG/DOSE (4 MG/3 ML), 2 MG/DOSE (8 MG/3 ML)
- TRULICITY
- RYBELSUS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Must meet all of the following 1.) The drug is prescribed for an FDA-approved indication, 2.) For a diagnosis of Type 2 Diabetes Mellitus the patient has a trial and failure, contraindication or intolerance to metformin
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No



# GLYCEROL PHENYLBUTYRATE

## Products Affected

- RAVICTI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	UREA CYCLE DISORDER (UCD): INITIAL: DIAGNOSIS IS CONFIRMED BY ENZYMATIC, BIOCHEMICAL OR GENETIC TESTING
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS
Other Criteria	UCD: INITIAL: TRIAL OF OR CONTRAINDICATION TO SODIUM PHENYLBUTYRATE. RENEWAL: PATIENT HAS CLINICAL BENEFIT FROM BASELINE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# GOSERELIN

## Products Affected

- ZOLADEX

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	ENDOMETRIOSIS: DIAGNOSIS IS CONFIRMED VIA SURGICAL OR DIRECT VISUALIZATION (E.G., PELVIC ULTRASOUND) OR HISTOPATHOLOGICAL CONFIRMATION (E.G., LAPAROSCOPY OR LAPAROTOMY) IN THE LAST 10 YEARS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	ENDOMETRIOSIS: PRESCRIBED BY OR IN CONSULTATION WITH AN OBSTETRICIAN/GYNECOLOGIST.
<b>Coverage Duration</b>	STAGE B2-C PROSTATIC CARCINOMA: 4 MOS. ENDOMETRIOSIS: 6 MOS PER LIFETIME. ALL OTHERS: 12 MONTHS.
<b>Other Criteria</b>	ENDOMETRIOSIS: 1) NO CONCURRENT USE WITH ANOTHER GNRH-MODULATING AGENT, 2) TRIAL OF OR CONTRAINDICATION TO NSAID AND PROGESTIN-CONTAINING PREPARATION, AND 3) HAS NOT RECEIVED A TOTAL OF 6 MONTHS OF TREATMENT PER LIFETIME. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# GUSELKUMAB

## Products Affected

- TREMFYA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE OR GENITAL AREA.
Age Restrictions	
Prescriber Restrictions	INITIAL: PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR DERMATOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	INITIAL: PSA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG). PSO: TRIAL OF OR CONTRAINDICATION ONE CONVENTIONAL THERAPY SUCH AS A PUVA (PHOTOTHERAPY ULTRAVIOLET LIGHT A), UVB (ULTRAVIOLET LIGHT B), TOPICAL CORTICOSTEROIDS, CALCIPOTRIENE, ACITRETIN, METHOTREXATE, OR CYCLOSPORINE. RENEWAL: PSO, PSA: CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# HIGH CONCENTRATION ORAL OPIOID SOLUTIONS

## Products Affected

- *morphine concentrate oral solution*
- *oxycodone oral concentrate*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	OPIOID TOLERANT: 12 MONTHS. HOSPICE, PALLIATIVE CARE OR END OF LIFE CARE: LIFETIME.
Other Criteria	1) OPIOID TOLERANT (I.E. PREVIOUS USE OF 60 MG ORAL MORPHINE PER DAY, 25 MCG TRANSDERMAL FENTANYL PER HOUR, 30 MG ORAL OXYCODONE PER DAY, 8 MG ORAL HYDROMORPHONE PER DAY, 25 MG ORAL OXYMORPHONE PER DAY, 60 MG ORAL HYDROCODONE PER DAY, OR AN EQUIANALGESIC DOSE OF ANOTHER OPIOID) AND HAS TROUBLE SWALLOWING OPIOID TABLETS, CAPSULES, OR LARGE VOLUMES OF LIQUID, OR 2) ENROLLED IN HOSPICE OR PALLIATIVE CARE OR END OF LIFE CARE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# HYDROXYUREA

---

## Products Affected

- SIKLOS ORAL TABLET 100 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# IBRUTINIB

---

## Products Affected

- IMBRUVICA ORAL CAPSULE 140 MG, 70 MG
- IMBRUVICA ORAL SUSPENSION
- IMBRUVICA ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# IBUPROFEN-FAMOTIDINE

## Products Affected

- *ibuprofen-famotidine*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	TRIAL OF ONE OF THE FOLLOWING GENERIC, FEDERAL LEGEND HISTAMINE H2-RECEPTOR ANTAGONISTS: FAMOTIDINE, CIMETIDINE, OR NIZATIDINE, AND TRIAL OF GENERIC, FEDERAL LEGEND IBUPROFEN.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# ICATIBANT

## Products Affected

- *icatibant*
- *sajazir*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	HEREDITARY ANGIOEDEMA (HAE): DIAGNOSIS CONFIRMED BY COMPLEMENT TESTING.
Age Restrictions	
Prescriber Restrictions	HAE: PRESCRIBED BY OR IN CONSULTATION WITH AN ALLERGIST, IMMUNOLOGIST, OR HEMATOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	HAE: NO CONCURRENT USE WITH OTHER MEDICATIONS FOR TREATMENT OF ACUTE HAE ATTACKS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# IDEALALISIB

---

## Products Affected

- ZYDELIG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# IMATINIB

## Products Affected

- *imatinib oral tablet 100 mg, 400 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	ADJUVANT GASTROINTESTINAL STROMAL TUMOR TREATMENT: 36 MONTHS. ALL OTHER DIAGNOSES: 12 MONTHS.
Other Criteria	PHILADELPHIA CHROMOSOME POSITIVE CHRONIC MYELOID LEUKEMIA: PATIENT HAS NOT RECEIVED A PREVIOUS TREATMENT WITH ANOTHER TYROSINE KINASE INHIBITOR.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# INFLIXIMAB

## Products Affected

- *infliximab*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, GENITAL AREA, OR FACE.
Age Restrictions	
Prescriber Restrictions	INITIAL: RHEUMATOID ARTHRITIS (RA), ANKYLOSING SPONDYLITIS (AS): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: RA: ONE OF THE FOLLOWING: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. PSA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, COSENTYX, ENBREL, XELJANZ, TREMFYA, RINVOQ, SKYRIZI. PSO: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, COSENTYX, ENBREL, SKYRIZI, TREMFYA. AS: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, COSENTYX, ENBREL, XELJANZ, RINVOQ. CD: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS, WHERE AGES ALIGN: HUMIRA, STELARA, SKYRIZI, RINVOQ. UC: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS, WHERE AGES ALIGN: HUMIRA, STELARA, XELJANZ, RINVOQ. RENEWAL: RA, AS, PSO, PSA: CONTINUES TO BENEFIT FROM THE MEDICATION. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# INFLIXIMAB-AXXQ

## Products Affected

- AVSOLA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, GENITAL AREA, OR FACE.
Age Restrictions	
Prescriber Restrictions	INITIAL: RHEUMATOID ARTHRITIS (RA), ANKYLOSING SPONDYLITIS (AS): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: RA: ONE OF THE FOLLOWING: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. PSA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, COSENTYX, ENBREL, XELJANZ, TREMFYA, RINVOQ, SKYRIZI. PSO: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, COSENTYX, ENBREL, SKYRIZI, TREMFYA. AS: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, COSENTYX, ENBREL, XELJANZ, RINVOQ. CD: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS, WHERE AGES ALIGN: HUMIRA, STELARA, SKYRIZI, RINVOQ. UC: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS, WHERE AGES ALIGN: HUMIRA, STELARA, XELJANZ, RINVOQ. RENEWAL: RA, AS, PSO, PSA: CONTINUES TO BENEFIT FROM THE MEDICATION. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# INTERFERON FOR MS-AVONEX

---

## Products Affected

- AVONEX INTRAMUSCULAR PEN  
INJECTOR KIT
- AVONEX INTRAMUSCULAR  
SYRINGE KIT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# INTERFERON FOR MS-BETASERON

---

## Products Affected

- BETASERON SUBCUTANEOUS KIT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# INTERFERON FOR MS-PLEGRIDY

## Products Affected

- PLEGRIDY SUBCUTANEOUS PEN INJECTOR 125 MCG/0.5 ML, 63 MCG/0.5 ML- 94 MCG/0.5 ML
- PLEGRIDY SUBCUTANEOUS SYRINGE 125 MCG/0.5 ML, 63 MCG/0.5 ML- 94 MCG/0.5 ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# INTERFERON GAMMA-1B

## Products Affected

- ACTIMMUNE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: CHRONIC GRANULOMATOUS DISEASE (CGD): PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST, INFECTIOUS DISEASE SPECIALIST, OR IMMUNOLOGIST. SEVERE MALIGNANT OSTEOPETROSIS (SMO): PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	RENEWAL: CGD, SMO: 1) DEMONSTRATED CLINICAL BENEFIT COMPARED TO BASELINE, AND 2) HAS NOT RECEIVED HEMATOPOIETIC CELL TRANSPLANTATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# ITRACONAZOLE SOLUTION

## Products Affected

- *itraconazole oral solution*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 MONTHS
Other Criteria	ESOPHAGEAL CANDIDIASIS AND OROPHARYNGEAL CANDIDIASIS; TRIAL OF OR CONTRAINDICATION TO FLUCONAZOLE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# IVACAFTOR

## Products Affected

- KALYDECO ORAL GRANULES IN PACKET 25 MG, 5.8 MG, 50 MG, 75 MG
- KALYDECO ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	CYSTIC FIBROSIS (CF): INITIAL: CONFIRMED MUTATION IN CFTR GENE ACCEPTABLE FOR THE TREATMENT OF CYSTIC FIBROSIS
Age Restrictions	
Prescriber Restrictions	CF: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST OR CYSTIC FIBROSIS EXPERT
Coverage Duration	INITIAL: 12 MONTHS. RENEWAL: LIFETIME
Other Criteria	CF: INITIAL: NOT HOMOZYGOUS FOR F508DEL MUTATION IN CFTR GENE. RENEWAL: 1) MAINTAINED, IMPROVED, OR DEMONSTRATED LESS THAN EXPECTED DECLINE IN FEV1 OR BODY MASS INDEX (BMI), OR 2) REDUCTION IN NUMBER OF PULMONARY EXACERBATIONS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# IVERMECTIN

## Products Affected

- *ivermectin oral*

PA Criteria	Criteria Details
Exclusion Criteria	Prevention or treatment of COVID-19
Required Medical Information	Diagnosis of one of the following: A.) Strongyloidiasis of the intestinal tract or B.) Onchocerciasis
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	1 month
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

# IVOSIDENIB

---

## Products Affected

- TIBSOVO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# IXAZOMIB

---

## Products Affected

- NINLARO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# IXEKIZUMAB

## Products Affected

- TALTZ AUTOINJECTOR
- TALTZ SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE, OR GENITAL AREA. NON-RADIOGRAPHIC AXIAL SPONDYLOARTHRITIS (NR-AXSPA): 1) C-REACTIVE PROTEIN (CRP) LEVELS ABOVE THE UPPER LIMIT OF NORMAL, OR 2) SACROILIITIS ON MAGNETIC RESONANCE IMAGING (MRI).
Age Restrictions	
Prescriber Restrictions	INITIAL: PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR DERMATOLOGIST. ANKYLOSING SPONDYLITIS (AS), NR-AXSPA: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.



PA Criteria	Criteria Details
Other Criteria	INITIAL: PSO: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS, WHERE AGES ALIGN: HUMIRA, COSENTYX, STELARA, ENBREL, SKYRIZI, TREMFYA. PSA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, COSENTYX, STELARA, ENBREL, XELJANZ, TREMFYA, RINVOQ, SKYRIZI. AS: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: ENBREL, HUMIRA, COSENTYX, XELJANZ, RINVOQ. NR-AXSPA: TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING PREFERRED AGENTS: COSENTYX, RINVOQ. RENEWAL: PSO, PSA, AS, NR-AXSPA: CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# LANADELUMAB

## Products Affected

- TAKHZYRO SUBCUTANEOUS (150 MG/ML) SOLUTION
- TAKHZYRO SUBCUTANEOUS SYRINGE 150 MG/ML, 300 MG/2 ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	HEREDITARY ANGIOEDEMA (HAE): INITIAL: DIAGNOSIS CONFIRMED BY COMPLEMENT TESTING.
Age Restrictions	
Prescriber Restrictions	HAE: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN ALLERGIST, IMMUNOLOGIST, OR HEMATOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	HAE: INITIAL: NOT ON CONCURRENT TREATMENT WITH ALTERNATIVE PROPHYLACTIC AGENT FOR HAE ATTACKS. RENEWAL: 1) IMPROVEMENT COMPARED TO BASELINE IN HAE ATTACKS (I.E., REDUCTIONS IN ATTACK FREQUENCY OR ATTACK SEVERITY), AND 2) NOT ON CONCURRENT TREATMENT WITH ALTERNATIVE PROPHYLACTIC AGENT FOR HAE ATTACKS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# LANREOTIDE

## Products Affected

- *lanreotide*
- SOMATULINE DEPOT  
SUBCUTANEOUS SYRINGE 120  
MG/0.5 ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	ACROMEGALY: INITIAL: THERAPY IS PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.
Coverage Duration	ACROMEGALY: INITIAL: 3 MOS, RENEWAL: 12 MOS.GEP-NETS, CARCINOID SYNDROME: 12 MOS.
Other Criteria	ACROMEGALY: INITIAL: TRIAL OF OR CONTRAINDICATION TO ONE GENERIC OCTREOTIDE INJECTION. RENEWAL: 1) REDUCTION, NORMALIZATION, OR MAINTENANCE OF IGF-1 LEVELS BASED ON AGE AND GENDER, AND 2) IMPROVEMENT OR SUSTAINED REMISSION OF CLINICAL SYMPTOMS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# LAPATINIB

---

## Products Affected

- *lapatinib*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# LAROTRECTINIB

## Products Affected

- VITRAKVI ORAL CAPSULE 100 MG, 25 MG
- VITRAKVI ORAL SOLUTION

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	APPROVAL FOR VITRAKVI ORAL SOLUTION: TRIAL OF VITRAKVI CAPSULES OR PATIENT IS UNABLE TO TAKE CAPSULE FORMULATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# LEDIPASVIR-SOFOSBUVIR

## Products Affected

- HARVONI ORAL PELLETS IN PACKET 33.75-150 MG, 45-200 MG
- HARVONI ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	HCV RNA LEVEL WITHIN PAST 6 MONTHS.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE.
Other Criteria	1) CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE, AND 2) NOT CONCURRENTLY TAKING ANY OF THE FOLLOWING: CARBAMAZEPINE, PHENYTOIN, PHENOBARBITAL, OXCARBAZEPINE, RIFAMPIN, RIFABUTIN, RIFAPENTINE, ROSUVASTATIN, TIPRANA VIR/RITONAVIR, SOFOSBUVIR (AS A SINGLE AGENT), EPCLUSA, ZEPATIER, MAVYRET, OR VOSEVI. REQUESTS FOR HARVONI 45MG-200MG PELLETS: PATIENT IS UNABLE TO SWALLOW TABLETS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# LENALIDOMIDE

---

## Products Affected

- *lenalidomide*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# LENVATINIB

---

## Products Affected

- LENVIMA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# LETERMOVIR

---

## Products Affected

- PREVYMIS ORAL

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	4 MONTHS
Other Criteria	NONE
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# LEUPROLIDE

---

## Products Affected

- *leuprolide subcutaneous kit*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	NONE
Coverage Duration	12 MONTHS
Other Criteria	BVD DETERMINATION AS REQUIRED PER CMS GUIDANCE
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

# LEUPROLIDE DEPOT

## Products Affected

- *leuprolide (3 month)*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# LEUPROLIDE-ELIGARD

---

## Products Affected

- ELIGARD
- ELIGARD (3 MONTH)
- ELIGARD (4 MONTH)
- ELIGARD (6 MONTH)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# LEUPROLIDE-LUPRON DEPOT

## Products Affected

- LUPRON DEPOT
- LUPRON DEPOT (3 MONTH)
- LUPRON DEPOT (4 MONTH)
- LUPRON DEPOT (6 MONTH)
- LUPRON DEPOT-PED  
INTRAMUSCULAR KIT 11.25 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: ENDOMETRIOSIS: DIAGNOSIS IS CONFIRMED VIA SURGICAL OR DIRECT VISUALIZATION (E.G., PELVIC ULTRASOUND) OR HISTOPATHOLOGICAL CONFIRMATION (E.G., LAPAROSCOPY OR LAPAROTOMY) IN THE LAST 10 YEARS.
Age Restrictions	
Prescriber Restrictions	INITIAL: ENDOMETRIOSIS: PRESCRIBED BY OR IN CONSULTATION WITH AN OBSTETRICIAN/GYNECOLOGIST.
Coverage Duration	PROSTATE CA: 12 MOS. UTERINE FIBROIDS: 3 MOS. ENDOMETRIOSIS: INITIAL/RENEWAL: 6 MOS.
Other Criteria	INITIAL: ENDOMETRIOSIS: 1) NO CONCURRENT USE WITH ANOTHER GNRH-MODULATING AGENT, 2) TRIAL OF OR CONTRAINDICATION TO NSAID AND PROGESTIN-CONTAINING PREPARATION, AND 3) HAS NOT RECEIVED A TOTAL OF 12 MONTHS OF TREATMENT PER LIFETIME. RENEWAL: ENDOMETRIOSIS: 1) IMPROVEMENT OF PAIN RELATED TO ENDOMETRIOSIS WHILE ON THERAPY, 2) RECEIVING CONCOMITANT ADD-BACK THERAPY (I.E., COMBINATION ESTROGEN-PROGESTIN OR PROGESTIN-ONLY CONTRACEPTIVE PREPARATION), 3) NO CONCURRENT USE WITH ANOTHER GNRH-MODULATING AGENT, AND 4) HAS NOT RECEIVED A TOTAL OF 12 MONTHS OF TREATMENT PER LIFETIME. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# LEVODOPA

## Products Affected

- INBRIJA INHALATION CAPSULE,  
W/INHALATION DEVICE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	PARKINSONS DISEASE (PD): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	PD: INITIAL: 1) NOT CURRENTLY TAKING MORE THAN 1600MG OF LEVODOPA PER DAY, AND 2) PHYSICIAN HAS OPTIMIZED DRUG THERAPY FOR PARKINSONS DISEASE. RENEWAL: IMPROVEMENT WITH MOTOR FLUCTUATIONS DURING OFF EPISODES WITH THE USE OF INBRIJA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# L-GLUTAMINE

## Products Affected

- ENDARI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	SICKLE CELL DISEASE(SCD): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST
Coverage Duration	INITIAL: 12 MONTHS. RENEWAL: LIFETIME.
Other Criteria	SCD: INITIAL: PATIENTS 18 YEARS OR OLDER: ONE OF THE FOLLOWING: 1) AT LEAST 2 SICKLE CELL CRISES IN THE PAST YEAR, 2) SICKLE-CELL ASSOCIATED SYMPTOMS WHICH ARE INTERFERING WITH ACTIVITIES OF DAILY LIVING, OR 3) HISTORY OF OR HAS RECURRENT ACUTE CHEST SYNDROME. PATIENTS 5 TO 17 YEARS: APPROVED WITHOUT ADDITIONAL CRITERIA. RENEWAL: HAS MAINTAINED OR EXPERIENCED REDUCTION IN ACUTE COMPLICATIONS OF SICKLE CELL DISEASE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# LIDOCAINE

## Products Affected

- *lidocaine hcl mucous membrane solution 4 % (40 mg/ml)*
- *lidocaine topical adhesive patch,medicated 5 %*
- *lidocaine topical ointment*
- **ZTLIDO**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

# LIDOCAINE PRILOCAINE

## Products Affected

- *lidocaine-prilocaine topical cream*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG MAY BE EITHER BUNDLED WITH AND COVERED UNDER END STAGE RENAL DISEASE DIALYSIS RELATED SERVICES OR COVERED UNDER MEDICARE D DEPENDING UPON THE CIRCUMSTANCES. INFORMATION MAY NEED TO BE SUBMITTED DESCRIBING THE USE AND SETTING OF THE DRUG TO MAKE THE DETERMINATION. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

# LOMITAPIDE

---

## Products Affected

- JUXTAPID ORAL CAPSULE 10 MG, 20 MG, 30 MG, 5 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST, ENDOCRINOLOGIST OR LIPIDOLOGIST.
Coverage Duration	12 MONTHS

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Other Criteria</b>	1) DIAGNOSIS DETERMINED BY A) DEFINITE SIMON BROOME DIAGNOSTIC CRITERIA, OR B) DUTCH LIPID NETWORK CRITERIA SCORE OF 8 OR GREATER, OR C) CLINICAL DIAGNOSIS BASED ON A HISTORY OF AN UNTREATED LDL-C CONCENTRATION GREATER THAN 500 MG/DL TOGETHER WITH EITHER XANTHOMA BEFORE 10 YEARS OF AGE, OR EVIDENCE OF HEFH IN BOTH PARENTS. 2) LDL-C LEVEL GREATER THAN OR EQUAL TO 70MG/DL WHILE ON MAXIMAL DRUG TREATMENT. 3) TRIAL OF EVOLOCUMAB UNLESS THE PATIENT HAS NON-FUNCTIONING LDL RECEPTORS. 4) MEETS ONE OF THE FOLLOWING: A) TAKING A HIGH-INTENSITY STATIN (I.E., ATORVASTATIN 40-80MG DAILY, ROSUVASTATIN 20-40MG DAILY) FOR A DURATION OF AT LEAST 8 WEEKS, B) TAKING A MAXIMALLY TOLERATED DOSE OF ANY STATIN FOR A DURATION OF AT LEAST 8 WEEKS GIVEN THAT THE PATIENT CANNOT TOLERATE A HIGH-INTENSITY STATIN, C) ABSOLUTE CONTRAINDICATION TO STATIN THERAPY (E.G., ACTIVE DECOMPENSATED LIVER DISEASE, NURSING FEMALE, PREGNANCY OR PLANS TO BECOME PREGNANT, HYPERSENSITIVITY REACTIONS), D) STATIN INTOLERANCE, OR E) TRIAL OF ROSUVASTATIN, ATORVASTATIN, OR STATIN THERAPY AT ANY DOSE AND HAS EXPERIENCED SKELETAL-MUSCLE RELATED SYMPTOMS (E.G., MYOPATHY).
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# LORLATINIB

---

## Products Affected

- LORBRENA ORAL TABLET 100 MG, 25 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# LUMACFTOR-IVACFTOR

## Products Affected

- ORKAMBI ORAL GRANULES IN PACKET
- ORKAMBI ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: CYSTIC FIBROSIS (CF): CONFIRMED MUTATION IN CFTR GENE ACCEPTABLE FOR THE TREATMENT OF CF.
Age Restrictions	
Prescriber Restrictions	CF: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST OR CF EXPERT.
Coverage Duration	INITIAL: 6 MONTHS, RENEWAL: LIFETIME.
Other Criteria	CF: RENEWAL: MAINTAINED, IMPROVED, OR DEMONSTRATED LESS THAN EXPECTED DECLINE IN FEV1 OR BODY MASS INDEX (BMI), OR REDUCTION IN NUMBER OF PULMONARY EXACERBATIONS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# MACITENTAN

## Products Affected

- OPSUMIT

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: 1) DOCUMENTED CONFIRMATORY DIAGNOSIS BASED ON RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: A) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, B) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, AND C) PULMONARY VASCULAR RESISTANCE (PVR) OF 3 WOOD UNITS OR GREATER, AND 2) NYHA-WHO FUNCTIONAL CLASS II-IV SYMPTOMS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	PAH: RENEWAL: 1) IMPROVEMENT FROM BASELINE IN THE 6-MINUTE WALK DISTANCE, OR 2) A STABLE 6-MINUTE WALK DISTANCE WITH A STABLE/IMPROVED WHO FUNCTIONAL CLASS.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# MEPOLIZUMAB

## Products Affected

- NUCALA SUBCUTANEOUS AUTO-INJECTOR
- NUCALA SUBCUTANEOUS RECON SOLN
- NUCALA SUBCUTANEOUS SYRINGE 100 MG/ML, 40 MG/0.4 ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: ASTHMA: BLOOD EOSINOPHIL LEVEL GREATER THAN OR EQUAL TO 150 CELLS/MCL WITHIN THE PAST 12 MONTHS.
Age Restrictions	
Prescriber Restrictions	INITIAL: ASTHMA: PRESCRIBED BY OR IN CONSULTATION WITH A PHYSICIAN SPECIALIZING IN PULMONARY OR ALLERGY MEDICINE. NASAL POLYPS: PRESCRIBED BY OR IN CONSULTATION WITH AN OTOLARYNGOLOGIST, ALLERGIST OR IMMUNOLOGIST.
Coverage Duration	INITIAL: ASTHMA: 4 MO. NASAL POLYPS: 6 MO. OTHERS: 12 MO. RENEWAL: NASAL POLYPS, ASTHMA: 12 MO.
Other Criteria	NONE
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# METHYLNALTREXONE

## Products Affected

- RELISTOR SUBCUTANEOUS SOLUTION
- RELISTOR SUBCUTANEOUS SYRINGE 12 MG/0.6 ML, 8 MG/0.4 ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	ADVANCED ILLNESS: OPIOID-INDUCED CONSTIPATION.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 MONTHS FOR PATIENTS RECEIVING PALLIATIVE CARE, 12 MONTHS FOR CHRONIC, NON-CANCER PAIN.
Other Criteria	ADVANCED ILLNESS: PATIENT IS RECEIVING PALLIATIVE CARE. CHRONIC NON-CANCER PAIN: PATIENT HAS BEEN TAKING OPIOIDS FOR AT LEAST 4 WEEKS AND HAD A PREVIOUS TRIAL OF OR CONTRAINDICATION TO NALOXEGOL (MOVANTIK) AND LUBIPROSTONE (AMITIZA).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# METHYLNALTREXONE ORAL

## Products Affected

- RELISTOR ORAL

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	OPIOID INDUCED CONSTIPATION WITH CHRONIC NON-CANCER PAIN: 1) HAS BEEN TAKING OPIOIDS FOR AT LEAST 4 WEEKS, AND 2) TRIAL OF OR CONTRAINDICATION TO NALOXEGOL (MOVANTIK) AND LUBIPROSTONE (AMITIZA).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# MIDOSTAURIN

---

## Products Affected

- RYDAPT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	ACUTE MYELOID LEUKEMIA: 6 MONTHS. ADVANCED SYSTEMIC MASTOCYTOSIS: 12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# MIFEPRISTONE

## Products Affected

- KORLYM

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	CUSHINGS SYNDROME (CS): INITIAL: DIAGNOSIS CONFIRMED BY ONE OF THE FOLLOWING: 1) 24-HR URINE FREE CORTISOL (2 OR MORE TESTS TO CONFIRM), 2) OVERNIGHT 1MG DEXAMETHASONE TEST, OR 3) LATE NIGHT SALIVARY CORTISOL (2 OR MORE TESTS TO CONFIRM).
Age Restrictions	
Prescriber Restrictions	CS: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.
Coverage Duration	INITIAL AND RENEWAL: 12 MONTHS.
Other Criteria	CS: INITIAL: HYPERCORTISOLISM IS NOT A RESULT OF CHRONIC GLUCOCORTICOIDS. RENEWAL: 1) CONTINUES TO HAVE IMPROVEMENT OF GLUCOSE TOLERANCE OR STABLE GLUCOSE TOLERANCE (E.G., REDUCED A1C, IMPROVED FASTING GLUCOSE, ETC.), 2) CONTINUES TO HAVE TOLERABILITY TO KORLYM, AND 3) CONTINUES TO NOT BE A CANDIDATE FOR SURGICAL TREATMENT OR HAS FAILED SURGERY.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# MIGALASTAT

## Products Affected

- GALAFOLD

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	FABRY DISEASE: INITIAL: PATIENT IS SYMPTOMATIC OR HAS EVIDENCE OF INJURY FROM GL-3 TO THE KIDNEY, HEART, OR CENTRAL NERVOUS SYSTEM RECOGNIZED BY LABORATORY, HISTOLOGICAL, OR IMAGING FINDINGS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	FABRY DISEASE: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEPHROLOGIST, CARDIOLOGIST, OR SPECIALIST IN GENETICS OR INHERITED METABOLIC DISORDERS.
<b>Coverage Duration</b>	INITIAL: 6 MOS. RENEWAL: 12 MOS.
<b>Other Criteria</b>	FABRY DISEASE: INITIAL: NOT CONCURRENTLY USING ENZYME REPLACEMENT THERAPY (I.E. FABRAZYME), RENEWAL: 1) PATIENT HAS DEMONSTRATED IMPROVEMENT OR STABILIZATION, AND 2) NOT CONCURRENTLY USING ENZYME REPLACEMENT THERAPY (I.E. FABRAZYME).
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# MIGLUSTAT

---

## Products Affected

- *miglustat*
- *yargesa*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# MILTEFOSINE

---

## Products Affected

- IMPAVIDO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# MOBOCERTINIB

---

## Products Affected

- EXKIVITY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# NAFARELIN

## Products Affected

- SYNAREL

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: ENDOMETRIOSIS: DIAGNOSIS IS CONFIRMED VIA SURGICAL OR DIRECT VISUALIZATION (E.G., PELVIC ULTRASOUND) OR HISTOPATHOLOGICAL CONFIRMATION (E.G., LAPAROSCOPY OR LAPAROTOMY) IN THE LAST 10 YEARS. CENTRAL PRECOCIOUS PUBERTY (CPP): FEMALES: ELEVATED LEVELS OF FOLLICLE-STIMULATING HORMONE (FSH) GREATER THAN 4.0 MIU/ML AND LUTEINIZING HORMONE (LH) LEVEL GREATER THAN 0.2 TO 0.3 MIU/ML AT DIAGNOSIS. MALES: ELEVATED LEVELS OF FSH GREATER THAN 5.0 MIU/ML AND LH LEVEL GREATER THAN 0.2 TO 0.3 MIU/ML AT DIAGNOSIS.
Age Restrictions	
Prescriber Restrictions	INITIAL: ENDOMETRIOSIS: PRESCRIBED BY OR IN CONSULTATION WITH AN OBSTETRICIAN/GYNECOLOGIST. CPP: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.
Coverage Duration	ENDOMETRIOSIS: 6 MONTHS. CPP: INITIAL/RENEWAL: 12 MONTHS.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Other Criteria</b>	<p>INITIAL: ENDOMETRIOSIS: 1) NO CONCURRENT USE WITH ANOTHER GNRH-MODULATING AGENT, 2) TRIAL OF OR CONTRAINDICATION TO NSAID AND PROGESTIN-CONTAINING PREPARATION, AND 3) HAS NOT RECEIVED A TOTAL OF 6 MONTHS OF TREATMENT PER LIFETIME.</p> <p>CPP: FEMALES: 1) YOUNGER THAN 8 YEARS OF AGE AT ONSET OF CPP, AND 2) DOCUMENTATION OF PUBERTAL STAGING USING THE TANNER SCALE FOR BREAST DEVELOPMENT (STAGE 2 OR ABOVE) AND PUBIC HAIR GROWTH (STAGE 2 OR ABOVE). MALES: 1) YOUNGER THAN 9 YEARS OF AGE AT ONSET OF CPP, AND 2) DOCUMENTATION OF PUBERTAL STAGING USING THE TANNER SCALE FOR GENITAL DEVELOPMENT (STAGE 2 OR ABOVE) AND PUBIC HAIR GROWTH (STAGE 2 OR ABOVE). RENEWAL: CPP: 1) TANNER SCALE STAGING AT INITIAL DIAGNOSIS OF CPP HAS STABILIZED OR REGRESSED DURING THREE SEPARATE MEDICAL VISITS IN THE PREVIOUS YEAR, AND 2) HAS NOT REACHED ACTUAL AGE WHICH CORRESPONDS TO CURRENT PUBERTAL AGE.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# NARCOLEPSY AGENTS

## Products Affected

- *armodafinil*
- *modafinil oral tablet 100 mg, 200 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

# NERATINIB MALEATE

## Products Affected

- NERLYNX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	EARLY-STAGE (STAGE I-III) BREAST CANCER: MEDICATION IS BEING REQUESTED WITHIN 2 YEARS OF COMPLETING THE LAST TRASTUZUMAB DOSE. ALL OTHER FDA APPROVED INDICATIONS ARE COVERED WITHOUT ADDITIONAL CRITERIA, EXCEPT THOSE CRITERIA IN THE FDA APPROVED LABEL.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# NILOTINIB

## Products Affected

- TASIGNA ORAL CAPSULE 150 MG, 200 MG, 50 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PREVIOUSLY TREATED CML: MUTATIONAL ANALYSIS PRIOR TO INITIATION AND TASIGNA IS APPROPRIATE PER NCCN GUIDELINE TABLE FOR TREATMENT RECOMMENDATIONS BASED ON BCR-ABL1 MUTATION PROFILE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# NINTEDANIB

## Products Affected

- OFEV

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of one of the following A.) Idiopathic pulmonary fibrosis (IPF), B.) Systemic sclerosis-associated interstitial lung disease (ILD), or C.) Chronic fibrosing interstitial lung disease with a progressive phenotype
Age Restrictions	18 years of age and older
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# NIRAPARIB

## Products Affected

- ZEJULA ORAL CAPSULE
- ZEJULA ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	RECURRENT EPITHELIAL OVARIAN, FALLOPIAN TUBE, OR PRIMARY PERITONEAL CANCER: 1) ZEJULA WILL BE USED AS MONOTHERAPY, AND 2) ZEJULA IS STARTED NO LATER THAN 8 WEEKS AFTER THE MOST RECENT PLATINUM-CONTAINING REGIMEN.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# NITISINONE

## Products Affected

- *nitisinone oral capsule 10 mg, 2 mg, 5 mg*
- ORFADIN ORAL CAPSULE 20 MG
- ORFADIN ORAL SUSPENSION

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	HEREDITARY TYROSINEMIA TYPE 1 (HT-1): INITIAL: DIAGNOSIS CONFIRMED BY ELEVATED URINARY OR PLASMA SUCCINYLACETONE LEVELS OR A MUTATION IN THE FUMARYLACETOACETATE HYDROLASE GENE. RENEWAL: URINARY OR PLASMA SUCCINYLACETONE LEVELS HAVE DECREASED FROM BASELINE WHILE ON TREATMENT WITH NITISINONE.
Age Restrictions	
Prescriber Restrictions	HT-1: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PRESCRIBER SPECIALIZING IN INHERITED METABOLIC DISEASES.
Coverage Duration	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.
Other Criteria	HT-1: INITIAL: ORFADIN SUSPENSION: TRIAL OF OR CONTRAINDICATION TO PREFERRED NITISINONE TABLETS OR CAPSULES.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# OBETICHOLIC ACID

## Products Affected

- OCALIVA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	PRIMARY BILIARY CHOLANGITIS (PBC): INITIAL/RENEWAL: COMPLETE BILIARY OBSTRUCTION.
<b>Required Medical Information</b>	PBC: INITIAL: DIAGNOSIS CONFIRMED BY TWO OF THE FOLLOWING: 1) ALKALINE PHOSPHATASE LEVEL OF AT LEAST 1.5 TIMES THE UPPER LIMIT OF NORMAL, 2) PRESENCE OF ANTIMITOCHONDRIAL ANTIBODIES AT A TITER OF 1:40 OR HIGHER, OR 3) HISTOLOGIC EVIDENCE OF NON-SUPPURATIVE DESTRUCTIVE CHOLANGITIS AND DESTRUCTION OF INTERLOBULAR BILE DUCTS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	PBC: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST OR HEPATOLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS
<b>Other Criteria</b>	PBC: INITIAL: USED IN COMBINATION WITH URSODEOXYCHOLIC ACID IN A PATIENT WITH AN INADEQUATE RESPONSE TO URSODEOXYCHOLIC ACID, OR AS MONOTHERAPY IN A PATIENT WHO IS UNABLE TO TOLERATE URSODEOXYCHOLIC ACID. RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# OFATUMUMAB-SQ

---

## Products Affected

- KESIMPTA PEN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# OJJAARA

## Products Affected

- OJJAARA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of intermediate or high-risk myelofibrosis (MF), including primary MF or secondary MF [postpolycythemia vera (PV) and post-essential thrombocythemia (ET)], in adults with anemia.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

# OLANZAPINE/SAMIDORPHAN

## Products Affected

- LYBALVI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	SCHIZOPHRENIA/BIPOLAR I: PRESCRIBED BY OR IN CONSULTATION WITH A PSYCHIATRIST
Coverage Duration	12 MONTHS
Other Criteria	SCHIZOPHRENIA: (1) PATIENT IS AT HIGH RISK OF WEIGHT GAIN AND (2) TRIAL OF OR CONTRAINDICATION TO LATUDA OR ONE OF THE FOLLOWING ORAL ANTIPSYCHOTICS: RISPERIDONE, CLOZAPINE TABLET, OLANZAPINE, IMMEDIATE RELEASE QUETIAPINE FUMARATE, ZIPRASIDONE, ARIPIPRAZOLE. BIPOLAR I: (1) PATIENT IS AT HIGH RISK OF WEIGHT GAIN AND (2) TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING ORAL ANTIPSYCHOTICS: RISPERIDONE, OLANZAPINE, IMMEDIATE RELEASE QUETIAPINE FUMARATE, ZIPRASIDONE, ARIPIPRAZOLE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# OLAPARIB

---

## Products Affected

- LYNPARZA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	NONE
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# OLUTASIDENIB

---

## Products Affected

- REZLIDHIA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# OMACETAXINE

---

## Products Affected

- SYNRIBO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# OMALIZUMAB

## Products Affected

- XOLAIR SUBCUTANEOUS RECON SOLN
- XOLAIR SUBCUTANEOUS SYRINGE 150 MG/ML, 75 MG/0.5 ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: ASTHMA: POSITIVE SKIN PRICK OR BLOOD TEST (E.G., ELISA, FEIA) TO A PERENNIAL AEROALLERGEN AND A BASELINE IGE SERUM LEVEL GREATER THAN OR EQUAL TO 30 IU/ML.
Age Restrictions	
Prescriber Restrictions	INITIAL AND RENEWAL: CHRONIC IDIOPATHIC URTICARIA (CIU): PRESCRIBED BY OR IN CONSULTATION WITH A PHYSICIAN SPECIALIZING IN ALLERGY OR PULMONARY MEDICINE, DERMATOLOGY OR IMMUNOLOGY. INITIAL: NASAL POLYPS: PRESCRIBED BY OR IN CONSULTATION WITH AN OTOLARYNGOLOGIST, ALLERGIST OR IMMUNOLOGIST. ASTHMA: PRESCRIBED BY OR IN CONSULTATION WITH A PHYSICIAN SPECIALIZING IN ALLERGY OR PULMONARY MEDICINE.
Coverage Duration	INITIAL: ASTHMA: 12 MO. CIU, NASAL POLYPS: 6 MO. RENEWAL: ASTHMA, NASAL POLYPS: 12 MO. CIU: 6 MO.



PA Criteria	Criteria Details
Other Criteria	<p>INITIAL: CIU: TRIAL OF OR CONTRAINDICATION TO A MAXIMALLY TOLERATED DOSE OF AN H1 ANTI-HISTAMINE AND STILL EXPERIENCES HIVES ON MOST DAYS OF THE WEEK. NASAL POLYPS: 1) PREVIOUS 90 DAY TRIAL OF ONE TOPICAL NASAL CORTICOSTEROID, AND 2) TRIAL OF OR CONTRAINDICATION TO ONE PREFERRED AGENT: NUCALA, DUPIXENT. ASTHMA: 1) CONCURRENT THERAPY WITH A MEDIUM, HIGH-DOSE OR MAXIMALLY TOLERATED DOSE OF AN INHALED CORTICOSTEROID AND AT LEAST ONE OTHER MAINTENANCE MEDICATION, 2) ONE OF THE FOLLOWING: (A) PATIENT EXPERIENCED AT LEAST ONE ASTHMA EXACERBATION REQUIRING SYSTEMIC CORTICOSTEROID BURST LASTING 3 OR MORE DAYS WITHIN THE PAST 12 MONTHS OR AT LEAST ONE SERIOUS EXACERBATION REQUIRING HOSPITALIZATION OR ER VISIT WITHIN THE PAST 12 MONTHS, OR (B) PATIENT HAS POOR SYMPTOM CONTROL DESPITE CURRENT THERAPY AS EVIDENCED BY AT LEAST THREE OF THE FOLLOWING WITHIN THE PAST 4 WEEKS: DAYTIME ASTHMA SYMPTOMS MORE THAN TWICE/WEEK, ANY NIGHT WAKING DUE TO ASTHMA, SABA RELIEVER FOR SYMPTOMS MORE THAN TWICE/WEEK, ANY ACTIVITY LIMITATION DUE TO ASTHMA, 3) NOT CONCURRENTLY RECEIVING DUPIXENT OR OTHER ANTI-IL5 BIOLOGICS WHEN THESE ARE USED FOR THE TREATMENT OF ASTHMA. RENEWAL: CIU: DIAGNOSIS OF CIU. NASAL POLYPS: CLINICAL BENEFIT COMPARED TO BASELINE. ASTHMA: 1) NOT CONCURRENTLY RECEIVING DUPIXENT OR OTHER ANTI-IL5 BIOLOGICS WHEN THESE ARE USED FOR THE TREATMENT OF ASTHMA, 2) CONTINUED USE OF ICS AND AT LEAST ONE OTHER MAINTENANCE MEDICATION, AND 3) CLINICAL RESPONSE AS EVIDENCED BY ONE OF THE FOLLOWING: A) REDUCTION IN ASTHMA</p>
	<p>EXACERBATIONS FROM BASELINE, B) DECREASED UTILIZATION OF RESCUE MEDICATIONS, C) REDUCTION IN SEVERITY OR FREQUENCY OF ASTHMA-RELATED SYMPTOMS, OR D) INCREASE IN PERCENT PREDICTED FEV1 FROM PRETREATMENT BASELINE.</p>
Indications	All FDA-approved Indications.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# OPICAPONE

---

## Products Affected

- ONGENTYS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	PARKINSONS DISEASE: 18 YEARS OF AGE OR OLDER
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# OSIMERTINIB

## Products Affected

- TAGRISSO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	EGFR EXON 19 DELETIONS OR EXON 21 L858R MUTATIONS NON-SMALL CELL LUNG CANCER (NSCLC) AND METASTATIC NSCLC WITH EGFR T790M MUTATION: NOT ON CONCURRENT THERAPY WITH AN EGFR TYROSINE KINASE-INHIBITOR.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# PACRITINIB

---

## Products Affected

- VONJO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	MYELOFIBROSIS: RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# PALBOCICLIB

---

## Products Affected

- IBRANCE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# PANOBINOSTAT

## Products Affected

- FARYDAK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS
Other Criteria	MULTIPLE MYELOMA: RENEWAL: TOLERATED THE FIRST 8 CYCLES OF THERAPY WITHOUT UNRESOLVED SEVERE OR MEDICALLY SIGNIFICANT TOXICITY.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# PARATHYROID HORMONE

## Products Affected

- NATPARA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	HYPOCALCEMIA SECONDARY TO HYPOPARATHYROIDISM: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	HYPOCALCEMIA SECONDARY TO HYPOPARATHYROIDISM: 1) TRIAL OF OR CONTRAINDICATION TO CALCITRIOL, 2) HYPOPARATHYROIDISM IS NOT DUE TO A CALCIUM SENSING RECEPTOR (CSR) MUTATION, AND 3) HYPOPARATHYROIDISM IS NOT CONSIDERED ACUTE POST-SURGICAL HYPOPARATHYROIDISM.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# PASIREOTIDE DIASPARTATE

## Products Affected

- SIGNIFOR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	CUSHINGS DISEASE (CD): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.
Other Criteria	CD: RENEWAL: 1) CONTINUED IMPROVEMENT OF CUSHINGS DISEASE, AND 2) MAINTAINED TOLERABILITY TO SIGNIFOR.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# PAZOPANIB

## Products Affected

- *pazopanib*
- VOTRIENT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	ADVANCED SOFT TISSUE SARCOMA (STS): NOT USED FOR ADIPOCYTIC STS OR GASTROINTESTINAL STROMAL TUMORS (GIST)
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# PDE5 INHIBITORS FOR PULMONARY ARTERIAL HYPERTENSION

## Products Affected

- *alyq*
- *sildenafil (pulm.hypertension) intravenous*
- *sildenafil (pulm.hypertension) oral tablet*
- *tadalafil (pulm. hypertension)*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	Any of the following A.) Nitrate therapy, including intermittent use, B.) Concomitant use with riociguat or other guanylate cyclase stimulators, C.) Concomitant use with HIV protease inhibitors or elvitegravir/cobicistat/tenofovir/emtricitabine
<b>Required Medical Information</b>	Diagnosis of pulmonary arterial hypertension (WHO Group I), confirmed by right heart catheterization or Doppler echocardiogram if patient is unable to undergo a right heart catheterization (e.g., patient is frail, elderly, etc.)
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST. THIS CRITERIA DOES NOT APPLY TO SILDENAFIL FOR AGES 1 TO 17 YEARS.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS
<b>Other Criteria</b>	NOT CONCURRENTLY OR INTERMITTENTLY TAKING ORAL ERECTILE DYSFUNCTION AGENTS (E.G. CIALIS, VIAGRA), ANY ORGANIC NITRATES IN ANY FORM, OR GUANYLATE CYCLASE STIMULATORS
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# PEGFILGRASTIM

---

## Products Affected

- FYLNETRA
- NEULASTA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	NON MYELOID MALIGNANCY, ACUTE RADIATION EXPOSURE: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	NON MYELOID MALIGNANCY: TRIAL OF OR CONTRAINDICATION TO NYVEPRIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# PEGFILGRASTIM - APGF

## Products Affected

- NYVEPRIA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	NON MYELOID MALIGNANCY: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# PEGFILGRASTIM - CBQV

## Products Affected

- UDENYCA
- UDENYCA AUTOINJECTOR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	NON MYELOID MALIGNANCY: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	NON MYELOID MALIGNANCY: TRIAL OF OR CONTRAINDICATION TO NYVEPRIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# PEGFILGRASTIM - JMDB

---

## Products Affected

- FULPHILA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	NON MYELOID MALIGNANCY: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	NON MYELOID MALIGNANCY: TRIAL OF OR CONTRAINDICATION TO NYVEPRIA
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# PEGFILGRASTIM-BMEZ

---

## Products Affected

- ZIEXTENZO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	NON MYELOID MALIGNANCY: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	NON MYELOID MALIGNANCY: TRIAL OF OR CONTRAINDICATION TO NYVEPRIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# PEGVALIASE-PQPZ

## Products Affected

- PALYNZIQ

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	PHENYLKETONURIA (PKU): INITIAL: NOT ON CONCURRENT TREATMENT WITH KUVAN. RENEWAL: 1) PATIENT CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NOT ON CONCURRENT TREATMENT WITH KUVAN .
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# PEGVISOMANT

---

## Products Affected

- SOMAVERT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# PEMIGATINIB

---

## Products Affected

- PEMAZYRE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# PENICILLAMINE

## Products Affected

- *penicillamine oral tablet*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: WILSONS DISEASE: CONFIRMED BY ONE OF THE FOLLOWING: 1) PLASMA COPPER-PROTEIN CERULOPLASMIN IS LESS THAN 20MG/DL, 2) LIVER BIOPSY POSITIVE FOR AN ABNORMALLY HIGH CONCENTRATION OF COPPER (GREATER THAN 250MCG/G DRY WEIGHT) OR THE PRESENCE OF KAYSER-FLEISCHER RINGS, OR 3) CONFIRMATION BY GENETIC TESTING FOR ATP7B MUTATIONS. CYSTINURIA: PATIENT HAS NEPHROLITHIASIS AND ONE OR MORE OF THE FOLLOWING: 1) STONE ANALYSIS SHOWING PRESENCE OF CYSTEINE, 2) IDENTIFICATION OF PATHOGNOMONIC HEXAGONAL CYSTINE CRYSTALS ON URINALYSIS, OR 3) POSITIVE FAMILY HISTORY OF CYSTINURIA WITH POSITIVE CYANIDE-NITROPRUSSIDE SCREEN.
Age Restrictions	
Prescriber Restrictions	WILSONS DISEASE: PRESCRIBED BY OR IN CONSULTATION WITH A HEPATOLOGIST OR GASTROENTEROLOGIST. CYSTINURIA: PRESCRIBED BY OR IN CONSULTATION WITH A NEPHROLOGIST. RHEUMATOID ARTHRITIS (RA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST.
Coverage Duration	INITIAL: 12 MONTHS, RENEWAL: LIFETIME.

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: WILSONS DISEASE: 1) KNOWN FAMILY HISTORY OF WILSONS DISEASE OR PHYSICAL EXAMINATION CONSISTENT WITH WILSONS DISEASE, AND 2) REQUESTS FOR FORMULARY VERSION OF PENICILLAMINE CAPSULE REQUIRE A TRIAL OF OR CONTRAINDICATION TO PENICILLAMINE TABLET (DEPEN). CYSTINURIA: REQUESTS FOR FORMULARY VERSION OF PENICILLAMINE CAPSULE REQUIRES A TRIAL OF OR CONTRAINDICATION TO PENICILLAMINE TABLET (DEPEN) AND A FORMULARY VERSION OF TIOPRONIN (THIOLA)/THIOLA EC. RA: 1) NO HISTORY OR OTHER EVIDENCE OF RENAL INSUFFICIENCY, 2) TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) - IF PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE GREATER THAN OR EQUAL TO 20 MG PER WEEK OR MAXIMALLY TOLERATED DOSE IS REQUIRED, AND 3) REQUESTS FOR FORMULARY VERSION OF PENICILLAMINE CAPSULE REQUIRES A TRIAL OF OR CONTRAINDICATION TO PENICILLAMINE TABLET (DEPEN). RENEWAL: RA: 1) NO HISTORY OR OTHER EVIDENCE OF RENAL INSUFFICIENCY, 2) EXPERIENCED OR MAINTAINED IMPROVEMENT IN TENDER JOINT COUNT OR SWOLLEN JOINT COUNT COMPARED TO BASELINE. WILSONS DISEASE, CYSTINURIA: PATIENT CONTINUES TO BENEFIT FROM THE MEDICATION.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# PEXIDARTINIB

---

## Products Affected

- TURALIO ORAL CAPSULE 200 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# PIMAVANSERIN

## Products Affected

- NUPLAZID

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	PSYCHOSIS IN PARKINSONS DISEASE (PD): INITIAL: 18 YEARS OR OLDER
Prescriber Restrictions	PSYCHOSIS IN PD: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST, GERIATRICIAN, OR A BEHAVIORAL HEALTH SPECIALIST (E.G., PSYCHIATRIST).
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS.
Other Criteria	PSYCHOSIS IN PD: RENEWAL: IMPROVEMENT IN PSYCHOSIS SYMPTOMS FROM BASELINE AND DEMONSTRATES A CONTINUED NEED FOR TREATMENT.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# PIRFENIDONE

## Products Affected

- *pirfenidone oral capsule*
- *pirfenidone oral tablet 267 mg, 534 mg, 801 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of idiopathic pulmonary fibrosis
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# PIRTOBRUTINIB

---

## Products Affected

- JAYPIRCA ORAL TABLET 100 MG, 50 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# POMALIDOMIDE

---

## Products Affected

- POMALYST

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# PONATINIB

## Products Affected

- ICLUSIG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	CML: MUTATIONAL ANALYSIS PRIOR TO INITIATION AND ICLUSIG IS APPROPRIATE PER NCCN GUIDELINE TABLE FOR TREATMENT RECOMMENDATIONS BASED ON BCR-ABL1 MUTATION PROFILE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# POSACONAZOLE

## Products Affected

- NOXAFIL ORAL SUSPENSION
- *posaconazole oral*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	OROPHARYNGEAL CANDIDIASIS (OPC): 3 MONTHS. PROPHYLAXIS: 6 MONTHS. TREATMENT: 12 WEEKS.
Other Criteria	NONE
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# POSACONAZOLE-POWDERMIX

## Products Affected

- NOXAFIL INTRAVENOUS
- NOXAFIL ORAL SUSP,DELAYED  
RELEASE FOR RECON

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 MONTHS
Other Criteria	PROPHYLAXIS OF INVASIVE ASPERGILLUS AND CANDIDA INFECTION: INABILITY TO SWALLOW TABLETS. CONTINUATION OF THERAPY AFTER HOSPITAL DISCHARGE REQUIRES NO EXTRA CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# PRALSETINIB

---

## Products Affected

- GAVRETO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# PRAMLINTIDE

---

## Products Affected

- SYMLINPEN 120
- SYMLINPEN 60

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	TYPE I OR TYPE II DIABETES: REQUIRING INSULIN OR CONTINUOUS INSULIN INFUSION (INSULIN PUMP) FOR GLYCEMIC CONTROL
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# PYRIMETHAMINE

## Products Affected

- *pyrimethamine*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	TOXOPLASMOSIS: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN INFECTIOUS DISEASE SPECIALIST.
Coverage Duration	TOXOPLASMOSIS: INITIAL: 8 WEEKS, RENEWAL: 6 MOS.
Other Criteria	TOXOPLASMOSIS: RENEWAL: ONE OF THE FOLLOWING: (1) PERSISTENT CLINICAL DISEASE (HEADACHE, NEUROLOGICAL SYMPTOMS, OR FEVER) AND PERSISTENT RADIOGRAPHIC DISEASE (ONE OR MORE MASS LESIONS ON BRAIN IMAGING), OR (2) CD4 COUNT LESS THAN 200 CELLS/MM3 AND CURRENTLY TAKING AN ANTI-RETROVIRAL THERAPY IF HIV POSITIVE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# QUININE

---

## Products Affected

- *quinine sulfate*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# REGORAFENIB

---

## Products Affected

- STIVARGA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# RELUGOLIX

---

## Products Affected

- ORGOVYX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# RIBOCICLIB

## Products Affected

- KISQALI FEMARA CO-PACK ORAL TABLET 200 MG/DAY(200 MG X 1)-2.5 MG, 400 MG/DAY(200 MG X 2)-2.5 MG, 600 MG/DAY(200 MG X 3)-2.5 MG
- KISQALI ORAL TABLET 200 MG/DAY (200 MG X 1), 400 MG/DAY (200 MG X 2), 600 MG/DAY (200 MG X 3)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	ADVANCED OR METASTATIC BREAST CANCER: TRIAL OF OR CONTRAINDICATION TO VERZENIO OR IBRANCE WHERE INDICATIONS ALIGN.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# RIFAXIMIN

---

## Products Affected

- XIFAXAN ORAL TABLET 200 MG, 550 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	TRAVELERS DIARRHEA/HE: 12 MOS. IBS-D: 8 WKS.
Other Criteria	RIFAXIMIN 550 MG TABLETS: HE: TRIAL OF OR CONTRAINDICATION TO LACTULOSE OR CONCURRENT LACTULOSE THERAPY.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# RIMEGEPANT

## Products Affected

- NURTEC ODT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	INITIAL: ACUTE MIGRAINE TREATMENT: TRIAL OF OR CONTRAINDICATION TO ONE TRIPTAN (E.G., SUMATRIPTAN, RIZATRIPTAN). EPISODIC MIGRAINE PREVENTION: 1) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION, AND 2) TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING PREVENTIVE MIGRAINE TREATMENTS: DIVALPROEX SODIUM, TOPIRAMATE, PROPRANOLOL, TIMOLOL. RENEWAL: ACUTE MIGRAINE TREATMENT: 1) IMPROVEMENT FROM BASELINE IN A VALIDATED ACUTE TREATMENT PATIENT-REPORTED OUTCOME QUESTIONNAIRE, OR 2) THERAPY WORKS CONSISTENTLY IN MAJORITY OF MIGRAINE ATTACKS. EPISODIC MIGRAINE PREVENTION: 1) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION, AND 2) REDUCTION IN MIGRAINE OR HEADACHE FREQUENCY, MIGRAINE SEVERITY, OR MIGRAINE DURATION WITH THERAPY.
Indications	All FDA-approved Indications.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# RIOCIGUAT

## Products Affected

- ADEMPAS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: PULMONARY ARTERIAL HYPERTENSION (PAH): 1) CONFIRMATORY DIAGNOSIS BASED ON RIGHT HEART CATHETERIZATION: A) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, B) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) 15 MMHG OR LESS, AND C) PULMONARY VASCULAR RESISTANCE (PVR) 3 WOOD UNITS OR GREATER, AND 2) NYHA-WHO FUNCTIONAL CLASS (FC) II-IV SYMPTOMS. PERSISTENT/RECURRENT CHRONIC THROMBOEMBOLIC PULMONARY HYPERTENSION (CTEPH) WHO GROUP 4: NYHA-WHO FUNCTIONAL CLASS II-IV SYMPTOMS.
Age Restrictions	
Prescriber Restrictions	INITIAL: PAH, CTEPH: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
Coverage Duration	INITIAL AND RENEWAL: 12 MONTHS.



<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Other Criteria</b>	INITIAL: PAH: NOT CONCURRENTLY TAKING NITRATES, NITRIC OXIDE DONORS, PHOSPHODIESTERASE INHIBITORS, OR NON-SPECIFIC PDE INHIBITORS. CTEPH: 1) NOT CONCURRENTLY TAKING NITRATES, NITRIC OXIDE DONORS, OR ANY PDE INHIBITORS, AND 2) NOT A CANDIDATE FOR SURGERY OR HAS INOPERABLE CTEPH OR HAS PERSISTENT OR RECURRENT DISEASE AFTER SURGICAL TREATMENT. RENEWAL: PAH, CTEPH: 1) IMPROVEMENT FROM BASELINE IN THE 6-MINUTE WALK DISTANCE OR A STABLE 6-MINUTE WALK DISTANCE WITH A STABLE/IMPROVED WHO FUNCTIONAL CLASS, AND 2) NOT CONCURRENTLY TAKING NITRATES, NITRIC OXIDE DONORS, PHOSPHODIESTERASE INHIBITORS, OR NON-SPECIFIC PDE INHIBITORS.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# RIPRETINIB

---

## Products Affected

- QINLOCK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# RISANKIZUMAB-RZAA

## Products Affected

- SKYRIZI INTRAVENOUS SYRINGE 150 MG/ML
- SKYRIZI SUBCUTANEOUS PEN INJECTOR • SKYRIZI SUBCUTANEOUS WEARABLE INJECTOR
- SKYRIZI SUBCUTANEOUS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: PLAQUE PSORIASIS (PSO): PLAQUE PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE OR GENITAL AREA.
Age Restrictions	
Prescriber Restrictions	INITIAL: PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR DERMATOLOGIST. CROHNS DISEASE (CD): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	INITIAL: PSO: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS PUVA (PHOTOTHERAPY ULTRAVIOLET LIGHT A), UVB (ULTRAVIOLET LIGHT B), TOPICAL CORTICOSTEROIDS, CALCIPOTRIENE, ACITRETIN, METHOTREXATE, OR CYCLOSPORINE. PSA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG). CD: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A CORTICOSTEROID (E.G., BUDESONIDE, METHYLPREDNISOLONE), AZATHIOPRINE, MERCAPTOPYRINE, METHOTREXATE, OR MESALAMINE. RENEWAL: PSO, PSA: CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# RISDIPLAM

## Products Affected

- EVRYSOI

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	SPINAL MUSCULAR ATROPHY (SMA): INITIAL: GENE MUTATION ANALYSIS INDICATING MUTATIONS OR DELETIONS OF BOTH ALLELES OF THE SURVIVAL MOTOR NEURON 1 (SMN1) GENE. FOR PRESYMPTOMATIC PATIENTS: UP TO THREE COPIES OF SURVIVAL MOTOR NEURON 2 (SMN2) BASED ON NEWBORN SCREENING.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	SMA: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROMUSCULAR SPECIALIST OR SMA SPECIALIST AT A SMA SPECIALTY CENTER.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS
<b>Other Criteria</b>	SMA: INITIAL: FOR SYMPTOMATIC PATIENTS: (1) BASELINE MOTOR FUNCTION ASSESSMENT BY A NEUROMUSCULAR SPECIALIST OR SMA SPECIALIST, AND (2) IF PATIENT RECEIVED GENE THERAPY, THE PATIENT HAD LESS THAN EXPECTED CLINICAL BENEFIT. RENEWAL: IMPROVED, MAINTAINED, OR DEMONSTRATED LESS THAN EXPECTED DECLINE IN: (1) MOTOR FUNCTION ASSESSMENTS COMPARED TO BASELINE, OR (2) OTHER MUSCLE FUNCTION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ROMOSOZUMAB

## Products Affected

- EVENITY SUBCUTANEOUS  
SYRINGE 210MG/2.34ML (  
105MG/1.17MLX2)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	POSTMENOPAUSAL OSTEOPOROSIS: 1) HAS NOT RECEIVED A TOTAL OF 12 MONTHS OF EVENITY TREATMENT, 2) ONE OF THE FOLLOWING: (A) HIGH RISK FOR FRACTURES DEFINED AS ONE OF THE FOLLOWING: (I) HISTORY OF OSTEOPOROTIC (I.E., FRAGILITY, LOW TRAUMA) FRACTURE(S), (II) 2 OR MORE RISK FACTORS FOR FRACTURE (E.G., HISTORY OF MULTIPLE RECENT LOW TRAUMA FRACTURES, BMD T-SCORE LESS THAN OR EQUAL TO -2.5, CORTICOSTEROID USE, USE OF GNRH ANALOGS SUCH AS NAFARELIN, ETC.), (III) NO PRIOR TREATMENT FOR OSTEOPOROSIS AND FRAX SCORE OF AT LEAST 20% FOR ANY MAJOR FRACTURE OR OF AT LEAST 3% FOR HIP FRACTURE, (B) UNABLE TO USE ORAL THERAPY (I.E., UPPER GASTROINTESTINAL PROBLEMS - UNABLE TO TOLERATE ORAL MEDICATION, LOWER GASTROINTESTINAL PROBLEMS - UNABLE TO ABSORB ORAL MEDICATIONS, TROUBLE REMEMBERING TO TAKE ORAL MEDICATIONS OR COORDINATING AN ORAL BISPHOSPHONATE WITH OTHER ORAL MEDICATIONS OR THEIR DAILY ROUTINE), OR (C) TRIAL OF, INTOLERANCE TO, OR A CONTRAINDICATION TO A BISPHOSPHONATE.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ROPEGINTERFERON ALFA-2B-NJFT

---

## Products Affected

- BESREMI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# RUCAPARIB

## Products Affected

- RUBRACA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	METASTATIC CASTRATION-RESISTANT PROSTATE CANCER: ONE OF THE FOLLOWING: 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# RUXOLITINIB

## Products Affected

- JAKAFI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	MYELOFIBROSIS: INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS. POLYCYTHEMIA VERA, GVHD: 12 MONTHS.
Other Criteria	MYELOFIBROSIS: RENEWAL: PATIENT CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# SAFINAMIDE

---

## Products Affected

- XADAGO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# SAPROPTERIN

## Products Affected

- *javygtor oral tablet, soluble*
- *sapropterin oral tablet, soluble*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL: 1 MONTH, RENEWAL 12 MONTHS.
Other Criteria	HYPERPHENYLALANINEMIA (HPA): INITIAL: NOT CONCURRENTLY USING PALYNZIQ. RENEWAL: 1) CONTINUES TO BENEFIT FROM TREATMENT, AND 2) NOT CONCURRENTLY USING PALYNZIQ.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# SARILUMAB

## Products Affected

- KEVZARA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	RHEUMATOID ARTHRITIS (RA): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST.
Coverage Duration	RA: INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS. POLYMYALGIA RHEUMATICA (PMR): 12 MONTHS.
Other Criteria	RA: INITIAL: ONE OF THE FOLLOWING: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# SATRALIZUMAB-MWGE

## Products Affected

- ENSPRYNG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	NMOSD: INITIAL: PRESCRIBED BY AN OPHTHALMOLOGIST OR PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS.
Other Criteria	NMOSD: INITIAL: 1) ONE OF THE FOLLOWING CORE CLINICAL CHARACTERISTIC: (A) OPTIC NEURITIS, (B) ACUTE MYELITIS, (C) AREA POSTREMA SYNDROME, (D) ACUTE BRAINSTEM SYNDROME, (E) SYMPTOMATIC NARCOLEPSY OR ACUTE DIENCEPHALIC CLINICAL SYNDROME WITH NMOSD-TYPICAL DIENCEPHALIC MRI LESIONS, OR (F) SYMPTOMATIC CEREBRAL SYNDROME WITH NMOSD-TYPICAL BRAIN LESIONS, AND 2) NO CONCURRENT USE WITH RITUXIMAB, INEBILIZUMAB, OR ECULIZUMAB. RENEWAL: 1) REDUCTION IN RELAPSE FREQUENCY FROM BASELINE, AND 2) NO CONCURRENT USE WITH USE RITUXIMAB, INEBILIZUMAB, OR ECULIZUMAB.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# SECUKINUMAB

## Products Affected

- COSENTYX (2 SYRINGES) SYRINGE 75 MG/0.5 ML
- COSENTYX INTRAVENOUS • COSENTYX UNOREADY PEN
- COSENTYX PEN (2 PENS)
- COSENTYX SUBCUTANEOUS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, GENITAL AREA, OR FACE. NON-RADIOGRAPHIC AXIAL SPONDYLOARTHRITIS (NR-AXSPA): 1) C-REACTIVE PROTEIN (CRP) LEVELS ABOVE THE UPPER LIMIT OF NORMAL, OR 2) SACROILIITIS ON MAGNETIC RESONANCE IMAGING (MRI).
Age Restrictions	
Prescriber Restrictions	INITIAL: PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR A DERMATOLOGIST. ANKYLOSING SPONDYLITIS (AS), NR-AXSPA, ENTHESITIS-RELATED ARTHRITIS (ERA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS

PA Criteria	Criteria Details
Other Criteria	INITIAL: PSO: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A PUVA (PHOTOTHERAPY ULTRAVIOLET LIGHT A), UVB (ULTRAVIOLET LIGHT B), TOPICAL CORTICOSTEROIDS, CALCIPOTRIENE, ACITRETIN, METHOTREXATE, OR CYCLOSPORINE. PSA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD (DISEASE-MODIFYING ANTI-RHEUMATIC DRUG). AS, NR-AXSPA: TRIAL OF OR CONTRAINDICATION TO AN NSAID (NON-STEROIDAL ANTI-INFLAMMATORY DRUG). ERA: TRIAL OF OR CONTRAINDICATION TO ONE NSAID, SULFASALAZINE, OR METHOTREXATE. RENEWAL: PSO, PSA, AS, NR-AXSPA, ERA: CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# SELEXIPAG

## Products Affected

- UPTRAVI ORAL TABLET 1,000 MCG, 1,200 MCG, 1,400 MCG, 1,600 MCG, 200 MCG, 400 MCG, 600 MCG, 800 MCG
- UPTRAVI ORAL TABLETS,DOSE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: 1) CONFIRMATORY DIAGNOSIS BASED ON RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: A) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, B) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) 15 MMHG OR LESS, AND C) PULMONARY VASCULAR RESISTANCE (PVR) 3 WOOD UNITS OR GREATER, AND 2) NYHA-WHO FUNCTIONAL CLASS (FC) II-IV SYMPTOMS.
Age Restrictions	
Prescriber Restrictions	PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Other Criteria</b>	PAH: INITIAL: WHO FC II-III SYMPTOMS: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING AGENTS, EACH FROM A DIFFERENT DRUG CLASS: 1) FORMULARY VERSION OF AN ORAL ENDOTHELIAL RECEPTOR ANTAGONIST, 2) FORMULARY VERSION OF AN ORAL PHOSPHODIESTERASE TYPE-5 INHIBITOR, OR 3) FORMULARY VERSION OF AN ORAL CGMP STIMULATOR. WHO FC III SYMPTOMS AND EVIDENCE OF RAPID PROGRESSION OR POOR PROGNOSIS, WHO FC IV SYMPTOMS: NO STEP. RENEWAL: 1) IMPROVEMENT FROM BASELINE IN THE 6-MINUTE WALK DISTANCE TEST, OR 2) REMAINED STABLE FROM BASELINE IN THE 6-MINUTE WALK DISTANCE TEST AND WHO FC HAS IMPROVED OR REMAINED STABLE.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# SELINEXOR

## Products Affected

- XPOVIO ORAL TABLET 100 MG/WEEK (50 MG X 2), 40 MG/WEEK (40 MG X 1), 40MG TWICE WEEK (40 MG X 2), 60 MG/WEEK (60 MG X 1), 60MG TWICE WEEK (120 MG/WEEK), 80 MG/WEEK (40 MG X 2), 80MG TWICE WEEK (160 MG/WEEK)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# SELPERCATINIB

---

## Products Affected

- RETEVMO ORAL CAPSULE 40 MG, 80 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# SELUMETINIB

---

## Products Affected

- KOSELUGO ORAL CAPSULE 10 MG, 25 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# SIPONIMOD

## Products Affected

- MAYZENT ORAL TABLET 0.25 MG, 1 MG, 2 MG
- MAYZENT STARTER(FOR 1MG MAINT)
- MAYZENT STARTER(FOR 2MG MAINT)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	MULTIPLE SCLEROSIS: RENEWAL: 1) DEMONSTRATION OF CLINICAL BENEFIT COMPARED TO PRE-TREATMENT BASELINE AND 2) DOES NOT HAVE LYMPHOPENIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# SODIUM OXYBATE

## Products Affected

- *sodium oxybate*
- XYREM

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: ALL INDICATIONS: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST, PSYCHIATRIST, OR SPECIALIST IN SLEEP MEDICINE.
Coverage Duration	INITIAL 6 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	INITIAL: EXCESSIVE DAYTIME SLEEPINESS (EDS) IN NARCOLEPSY: 1) NOT CURRENTLY TAKING A SEDATIVE HYPNOTIC AGENT, 2) FOR PATIENTS 18 YEARS OR OLDER: TRIAL OF OR CONTRAINDICATION TO THE FORMULARY VERSION OF MODAFINIL, ARMODAFINIL, PITOLISANT OR SOLRIAMFETOL AND ONE OTHER GENERIC STIMULANT INDICATED FOR EDS IN NARCOLEPSY, AND 3) FOR PATIENTS 7 TO 17 YEARS OF AGE: TRIAL OF OR CONTRAINDICATION TO ONE OTHER GENERIC STIMULANT INDICATED FOR EDS IN NARCOLEPSY. CATAPLEXY IN NARCOLEPSY: NOT CURRENTLY TAKING A SEDATIVE HYPNOTIC AGENT. RENEWAL (ALL INDICATIONS): 1) SUSTAINED IMPROVEMENT OF SYMPTOMS COMPARED TO BASELINE, AND 2) NOT CURRENTLY TAKING A SEDATIVE HYPNOTIC AGENT.
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No



# SOFOSBUVIR/VELPATASVIR

## Products Affected

- EPCLUSA ORAL PELLETS IN PACKET 150-37.5 MG, 200-50 MG
- EPCLUSA ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	HCV RNA LEVEL WITHIN PAST 6 MONTHS.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE.
Other Criteria	1) CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE, 2) NOT CONCURRENTLY TAKING ANY OF THE FOLLOWING MEDICATIONS: AMIODARONE, CARBAMAZEPINE, PHENYTOIN, PHENOBARBITAL, OXCARBAZEPINE, RIFAMPIN, RIFABUTIN, RIFAPENTINE, HIV REGIMEN THAT CONTAINS EFAVIRENZ, ROSUVASTATIN AT DOSES ABOVE 10MG, TIPRANA VIR/RITONAVIR, TOPOTECAN, SOVALDI (AS A SINGLE AGENT), HARVONI, ZEPATIER, MAVYRET, OR VOSEVI, AND 3) PATIENTS WITH DECOMPENSATED CIRRHOSIS REQUIRE CONCURRENT RIBAVIRIN UNLESS RIBAVIRIN INELIGIBLE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# SOFOSBUVIR/VELPATASVIR/VOXILAPREVIR

## Products Affected

- VOSEVI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	HCV RNA LEVEL WITHIN PAST 6 MONTHS
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE.
Other Criteria	1) CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE, 2) NOT CONCURRENTLY TAKING ANY OF THE FOLLOWING MEDICATIONS: AMIODARONE, CARBAMAZEPINE, PHENYTOIN, PHENOBARBITAL, OXCARBAZEPINE, RIFAMPIN, RIFABUTIN, RIFAPENTINE, CYCLOSPORINE, PITAVASTATIN, PRAVASTATIN (DOSES ABOVE 40MG), ROSUVASTATIN, METHOTREXATE, MITOXANTRONE, IMATINIB, IRINOTECAN, LAPATINIB, SULFASALAZINE, TOPOTECAN, OR HIV REGIMEN THAT CONTAINS EFAVIRENZ, ATAZANAVIR, LOPINAVIR, TIPRANAVIR/RITONAVIR, SOVALDI (AS A SINGLE AGENT), EPCLUSA, HARVONI, ZEPATIER, OR MAVYRET, AND 3) DOES NOT HAVE MODERATE OR SEVERE HEPATIC IMPAIRMENT (CHILD-PUGH B OR C).
Indications	All FDA-approved Indications.
Off Label Uses	

PA Criteria	Criteria Details
Part B Prerequisite	No

# SOLRIAMFETOL

## Products Affected

- SUNOSI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: EXCESSIVE DAYTIME SLEEPINESS (EDS) IN NARCOLEPSY: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST, PSYCHIATRIST, OR SPECIALIST IN SLEEP MEDICINE.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	INITIAL: EDS IN NARCOLEPSY: TRIED THE FORMULARY VERSION OF MODAFINIL OR ARMODAFINIL, AND ONE OTHER GENERIC STIMULANT INDICATED FOR EDS IN NARCOLEPSY. EDS IN OBSTRUCTIVE SLEEP APNEA (OSA): TRIED THE FORMULARY VERSION OF MODAFINIL OR ARMODAFINIL. RENEWAL: EDS IN NARCOLEPSY OR OSA: SUSTAINED IMPROVEMENT OF SYMPTOMS COMPARED TO BASELINE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# SOMATROPIN - NORDITROPIN

## Products Affected

- NORDITROPIN FLEXPRO

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	INITIAL/RENEWAL: ATHLETIC ENHANCEMENT, ANTI-AGING PURPOSES.
<b>Required Medical Information</b>	INITIAL: PEDIATRIC GROWTH HORMONE DEFICIENCY (GHD), IDIOPATHIC SHORT STATURE (ISS), SMALL FOR GESTATIONAL AGE (SGA), TURNER SYNDROME (TS), NOONAN SYNDROME: HEIGHT AT LEAST 2 STANDARD DEVIATIONS (SD) BELOW THE MEAN HEIGHT FOR NORMAL CHILDREN OF THE SAME AGE AND GENDER. PRADER WILLI SYNDROME (PWS): CONFIRMED GENETIC DIAGNOSIS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL/RENEWAL: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Other Criteria</b>	INITIAL: ADULT GHD: GROWTH HORMONE DEFICIENCY ALONE OR ASSOCIATED WITH MULTIPLE HORMONE DEFICIENCIES (HYPOPITUITARISM), AS A RESULT OF PITUITARY DISEASES, HYPOTHALAMIC DISEASE, SURGERY, RADIATION THERAPY, TRAUMA, OR CONTINUATION OF THERAPY FROM CHILDHOOD ONSET GROWTH HORMONE DEFICIENCY. PEDIATRIC GHD, ISS, SGA, TS, NOONAN SYNDROME: OPEN EPIPHYSES AS CONFIRMED BY RADIOGRAPH OF THE WRIST AND HAND. RENEWAL: PEDIATRIC GHD: 1) IMPROVEMENT WHILE ON THERAPY (I.E., INCREASED HEIGHT OR INCREASED GROWTH VELOCITY), AND 2) OPEN EPIPHYSES AS CONFIRMED BY RADIOGRAPH OF THE WRIST AND HAND OR PATIENT HAS NOT COMPLETED PREPUBERTALGROWTH. ISS, SGA, TS, NOONAN SYNDROME: 1) IMPROVEMENT WHILE ON THERAPY (I.E., INCREASED HEIGHT OR INCREASED GROWTH VELOCITY), AND 2) OPEN EPIPHYSES AS CONFIRMED BY RADIOGRAPH OF THE WRIST AND HAND. PWS: IMPROVEMENT IN BODY COMPOSITION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# SOMATROPIN - SEROSTIM

## Products Affected

- SAIZEN SUBCUTANEOUS RECON SOLN 5 MG
- SEROSTIM SUBCUTANEOUS RECON SOLN 4 MG, 5 MG, 6 MG
- ZOMACTON SUBCUTANEOUS RECON SOLN 5 MG

PA Criteria	Criteria Details
Exclusion Criteria	INITIAL/RENEWAL: ATHLETIC ENHANCEMENT, ANTI-AGING PURPOSES
Required Medical Information	INITIAL: HIV/WASTING: MEETS ONE OF THE FOLLOWING CRITERIA FOR WEIGHT LOSS: 1) 10% UNINTENTIONAL WEIGHT LOSS OVER 12 MONTHS, 2) 7.5% UNINTENTIONAL WEIGHT LOSS OVER 6 MONTHS, 3) 5% BODY CELL MASS (BCM) LOSS WITHIN 6 MONTHS, 4) BODY CELL MASS (BCM) LESS THAN 35% (MEN) OF TOTAL BODY WEIGHT AND BODY MASS INDEX (BMI) LESS THAN 27 KG PER METER SQUARED, 5) BCM LESS THAN 23% (WOMEN) OF TOTAL BODY WEIGHT AND A BODY MASS INDEX (BMI) LESS THAN 27 KG PER METER SQUARED, OR 6) BMI LESS THAN 18.5 KG PER METER SQUARED.
Age Restrictions	
Prescriber Restrictions	HIV/WASTING: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST, NUTRITIONAL SUPPORT SPECIALIST, OR INFECTIOUS DISEASE SPECIALIST.
Coverage Duration	INITIAL/RENEWAL: 3 MONTHS.
Other Criteria	HIV/WASTING: INITIAL: 1) CURRENTLY ON HIV ANTIRETROVIRAL THERAPY, AND 2) INADEQUATE RESPONSE TO ONE PREVIOUS THERAPY (E.G., MEGACE, APPETITE STIMULANTS, ANABOLIC STEROIDS). RENEWAL: 1) CURRENTLY ON HIV ANTIRETROVIRAL THERAPY, AND 2) CLINICAL BENEFIT IN MUSCLE MASS AND WEIGHT.
Indications	All FDA-approved Indications.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# SONIDEGIB

## Products Affected

- ODOMZO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	LOCALLY ADVANCED BASAL CELL CARCINOMA (BCC): BASELINE SERUM CREATINE KINASE (CK) AND SERUM CREATININE LEVELS
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# SORAFENIB

## Products Affected

- *sorafenib*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# SOTORASIB

## Products Affected

- LUMAKRAS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# STIRIPENTOL

## Products Affected

- DIACOMIT ORAL CAPSULE 250 MG, 500 MG
- DIACOMIT ORAL POWDER IN PACKET 250 MG, 500 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	DRAVET SYNDROME: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# SUNITINIB

## Products Affected

- *sunitinib malate*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	GASTROINTESTINAL STROMAL TUMORS (GIST): TRIAL OF OR CONTRAINDICATION TO IMATINIB (GLEEVEC).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# TADALAFIL

## Products Affected

- *tadalafil oral tablet 2.5 mg, 5 mg*

PA Criteria	Criteria Details
Exclusion Criteria	ERECTILE DYSFUNCTION WITHOUT DIAGNOSIS OF BENIGN PROSTATIC HYPERPLASIA (BPH).
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	BPH: 1) TRIAL OF ONE ALPHA BLOCKER (SUCH AS DOXAZOSIN, TERAZOSIN, TAMSULOSIN, ALFUZOSIN), AND 2) TRIAL OF ONE 5-ALPHA-REDUCTASE INHIBITOR (SUCH AS FINASTERIDE, DUTASTERIDE). APPLIES TO 2.5MG AND 5MG STRENGTHS ONLY. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# TALAZOPARIB

---

## Products Affected

- TALZENNA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	NONE
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# TASIMELTEON

## Products Affected

- HETLIOZ
- HETLIOZ LQ
- *tasimelteon*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	LIFETIME
Other Criteria	NON-24 HOUR SLEEP-WAKE DISORDER: PATIENT IS LIGHT-INSENSITIVE OR HAS TOTAL BLINDNESS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# TAZEMETOSTAT

---

## Products Affected

- TAZVERIK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# TBO-FILGRASTIM

## Products Affected

- GRANIX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	NON-MYELOID MALIGNANCIES: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	NON-MYELOID MALIGNANCIES: TRIAL OF OR CONTRAINDICATION TO THE PREFERRED AGENT: NIVESTYM.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# TEDUGLUTIDE

## Products Affected

- GATTEX 30-VIAL

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	SHORT BOWEL SYNDROME (SBS): INITIAL/RENEWAL: PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST
Coverage Duration	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS
Other Criteria	SBS: INITIAL: PATIENT IS DEPENDENT ON INTRAVENOUS PARENTERAL NUTRITION DEFINED AS REQUIRING PARENTERAL NUTRITION AT LEAST THREE TIMES PER WEEK. RENEWAL: ACHIEVED OR MAINTAINED A DECREASED NEED FOR PARENTERAL SUPPORT COMPARED TO BASELINE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# TELOTRISTAT

## Products Affected

- XERMELO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	CARCINOID SYNDROME DIARRHEA: PRESCRIBED BY OR IN CONSULTATION WITH AN ONCOLOGIST OR GASTROENTEROLOGIST
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# TEPOTINIB

---

## Products Affected

- TEPMETKO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# TERIFLUNOMIDE

---

## Products Affected

- AUBAGIO
- *teriflunomide*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# TESAMORELIN

---

## Products Affected

- EGRIFTA SV

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	3 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# TESTOSTERONE

## Products Affected

- XYOSTED

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: MALE HYPOGONADISM: CONFIRMED BY: 1) AT LEAST TWO TOTAL SERUM TESTOSTERONE LEVELS OF LESS THAN 300 NG/DL TAKEN ON SEPARATE OCCASIONS, OR 2) FREE SERUM TESTOSTERONE LEVEL OF LESS THAN 5 NG/DL.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	MALE HYPOGONADISM: INITIAL/RENEWAL:12 MO. ALL OTHER INDICATIONS: LIFETIME OF MEMBERSHIP IN PLAN.
Other Criteria	RENEWAL: MALE HYPOGONADISM: IMPROVED SYMPTOMS COMPARED TO BASELINE AND TOLERANCE TO TREATMENT.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# TETRABENAZINE

## Products Affected

- *tetrabenazine*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	HUNTINGTONS DISEASE: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST OR MOVEMENT DISORDER SPECIALIST
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# TEZACAFTOR/IVACAFTOR

## Products Affected

- SYMDEKO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	CYSTIC FIBROSIS (CF): INITIAL: CONFIRMED MUTATION IN CFTR GENE ACCEPTABLE FOR THE TREATMENT OF CYSTIC FIBROSIS.
Age Restrictions	
Prescriber Restrictions	CF: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST OR CYSTIC FIBROSIS EXPERT
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: LIFETIME
Other Criteria	CF: RENEWAL: MAINTAINED, IMPROVED, OR DEMONSTRATED LESS THAN EXPECTED DECLINE IN FEV1 OR BODY MASS INDEX (BMI), OR REDUCTION IN NUMBER OF PULMONARY EXACERBATIONS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# THALIDOMIDE

---

## Products Affected

- THALOMID

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# TILDRAKIZUMAB-ASMN

## Products Affected

- ILUMYA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	PLAQUE PSORIASIS (PSO): INITIAL: PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, GENITAL AREA, OR FACE.
Age Restrictions	
Prescriber Restrictions	PSO: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.
Other Criteria	PSO: INITIAL: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, COSENTYX, STELARA, ENBREL, SKYRIZI, TREMFYA. RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# TIVOZANIB

---

## Products Affected

- FOTIVDA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# TOCILIZUMAB IV

## Products Affected

- ACTEMRA

PA Criteria	Criteria Details
Exclusion Criteria	CORONAVIRUS DISEASE 2019 (COVID-19) IN HOSPITALIZED ADULTS
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: RHEUMATOID ARTHRITIS (RA), POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. SYSTEMIC JUVENILE IDIOPATHIC ARTHRITIS (SJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST, DERMATOLOGIST, OR IMMUNOLOGIST.
Coverage Duration	INITIAL: RA, PJIA, SJIA, GCA: 6 MONTHS. CRS: 1 MONTH. RENEWAL: RA, PJIA, SJIA, GCA: 12 MONTHS.
Other Criteria	INITIAL: RA: ONE OF THE FOLLOWING: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. PJIA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ IR. SJIA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG). RENEWAL: RA, PJIA, SJIA: CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TOCILIZUMAB SQ

## Products Affected

- ACTEMRA
- ACTEMRA ACTPEN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: RHEUMATOID ARTHRITIS (RA), POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. SYSTEMIC JUVENILE IDIOPATHIC ARTHRITIS (SJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST, DERMATOLOGIST, OR IMMUNOLOGIST. SYSTEMIC SCLEROSIS-ASSOCIATED INTERSTITIAL LUNG DISEASE (SSC-ILD): PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST OR RHEUMATOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.



PA Criteria	Criteria Details
Other Criteria	<p>INITIAL: RA: ONE OF THE FOLLOWING: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. PJIA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ IR. SJIA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG). SSC-ILD: DOES NOT HAVE OTHER KNOWN CAUSES OF INTERSTITIAL LUNG DISEASE (E.G., CONNECTIVE TISSUE DISEASE, DRUG TOXICITY, ASBESTOS OR BERYLLIUM EXPOSURE, HYPERSENSITIVITY PNEUMONITIS). RENEWAL: RA, PJIA, SJIA: CONTINUES TO BENEFIT FROM THE MEDICATION. SSC-ILD: CLINICAL MEANINGFUL IMPROVEMENT OR MAINTENANCE IN ANNUAL RATE OF DECLINE.</p>
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# TOFACITINIB

## Products Affected

- XELJANZ
- XELJANZ XR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: RHEUMATOID ARTHRITIS (RA), ANKYLOSING SPONDYLITIS (AS), POLYARTICULAR COURSE JUVENILE IDIOPATHIC ARTHRITIS (PCJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR DERMATOLOGIST. ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) - IF A PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE GREATER THAN OR EQUAL TO 20MG PER WEEK OR MAXIMALLY TOLERATED DOSE IS REQUIRED. PSA, PCJIA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD. AS: TRIAL OF OR CONTRAINDICATION TO AN NSAID. UC: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A CORTICOSTEROID (E.G., BUDESONIDE, METHYLPREDNISOLONE), AZATHIOPRINE, MERCAPTOPURINE, METHOTREXATE, OR MESALAMINE. RENEWAL: RA, PSA, AS, PCJIA: CONTINUES TO BENEFIT FROM THE MEDICATION.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TOLVAPTAN

## Products Affected

- JYNARQUE ORAL TABLET
- JYNARQUE ORAL TABLETS, SEQUENTIAL

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	AUTOSOMAL DOMINANT POLYCYSTIC KIDNEY DISEASE (ADPKD): INITIAL: 1) CONFIRMED POLYCYSTIC KIDNEY DISEASE VIA CT, MRI, OR ULTRASOUND, AND 2) GENETIC TESTING FOR CAUSATIVE MUTATIONS OR FAMILY HISTORY OF CONFIRMED POLYCYSTIC KIDNEY DISEASE IN ONE OR BOTH PARENTS.
Age Restrictions	
Prescriber Restrictions	ADPKD: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEPHROLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	ADPKD: INITIAL: DOES NOT HAVE ESRD (I.E., RECEIVING DIALYSIS OR HAS UNDERGONE RENAL TRANSPLANT). RENEWAL: HAS NOT PROGRESSED TO ESRD/DIALYSIS OR TRANSPLANT.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# TOPICAL TRETINOIN

## Products Affected

- ALTRENO
- *avita topical gel*
- *tretinoin*

PA Criteria	Criteria Details
Exclusion Criteria	COSMETIC INDICATIONS SUCH AS WRINKLES, PHOTOAGING, MELASMA.
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	ACNE VULGARIS: BRAND TOPICAL TRETINOIN REQUIRES TRIAL OF OR CONTRAINDICATION TO A GENERIC TOPICAL TRETINOIN PRODUCT.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# TRAMETINIB

---

## Products Affected

- MEKINIST ORAL TABLET 0.5 MG, 2 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# TREPROSTINIL INHALED

## Products Affected

- TYVASO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: PULMONARY ARTERIAL HYPERTENSION (PAH): (1) CONFIRMATORY DIAGNOSIS BASED ON RIGHT HEART CATHETERIZATION: A) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, B) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) 15 MMHG OR LESS, AND C) PULMONARY VASCULAR RESISTANCE (PVR) 3 WOOD UNITS OR GREATER, AND (2) NYHA-WHO FUNCTIONAL CLASS (FC) III-IV SYMPTOMS. PULMONARY HYPERTENSION-INTERSTITIAL LUNG DISEASE (PH-ILD): CONFIRMATORY DIAGNOSIS BASED ON RIGHT HEART CATHETERIZATION: (1) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, (2) PULMONARY VASCULAR RESISTANCE (PVR) 3 WOOD UNITS OR GREATER, AND (3) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS.
Age Restrictions	
Prescriber Restrictions	INITIAL: PAH, PH-ILD: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
Coverage Duration	INITIAL: PAH: 12 MONTHS, PH-ILD: 6 MONTHS. RENEWAL: PAH, PH-ILD: 12 MONTHS.

PA Criteria	Criteria Details
Other Criteria	<p>INITIAL: PAH: WHO FC III SYMPTOMS: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING AGENTS FROM DIFFERENT DRUG CLASSES: 1) FORMULARY VERSION OF AN ORAL ENDOTHELIAL RECEPTOR ANTAGONIST, 2) FORMULARY VERSION OF AN ORAL PHOSPHODIESTERASE TYPE-5 INHIBITOR, OR 3) FORMULARY VERSION OF AN ORAL CGMP STIMULATOR. WHO FC III SYMPTOMS AND EVIDENCE OF RAPID PROGRESSION OR POOR PROGNOSIS, WHO FC IV SYMPTOMS: TRIAL OF OR CONTRAINDICATION TO A FORMULARY VERSION OF AN IV/SQ PROSTACYCLIN. RENEWAL: PAH: IMPROVEMENT FROM BASELINE IN THE 6-MINUTE WALK DISTANCE TEST OR REMAINED STABLE FROM BASELINE IN THE 6-MINUTE WALK DISTANCE TEST AND WHO FC HAS IMPROVED OR REMAINED STABLE. PH-ILD: IMPROVEMENT FROM BASELINE IN THE 6-MINUTE WALK DISTANCE TEST OR A STABLE 6-MINUTE WALK DISTANCE TEST. THIS DRUG MAY BE COVERED UNDER MEDICARE PART B OR D DEPENDING UPON THE CIRCUMSTANCES. INFORMATION MAY NEED TO BE SUBMITTED DESCRIBING THE USE AND SETTING OF THE DRUG TO MAKE THE DETERMINATION.</p>
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# TREPROSTINIL INJECTABLE

## Products Affected

- *treprostinil sodium*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: (1) CONFIRMATORY DIAGNOSIS BASED ON RIGHT HEART CATHETERIZATION: A) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, B) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) 15 MMHG OR LESS, AND C) PULMONARY VASCULAR RESISTANCE (PVR) 3 WOOD UNITS OR GREATER, AND (2) NYHA-WHO FUNCTIONAL CLASS (FC) II-IV SYMPTOMS.
Age Restrictions	
Prescriber Restrictions	PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
Coverage Duration	PAH: INITIAL AND RENEWAL: 12 MONTHS
Other Criteria	PAH: INITIAL: CONTINUATION FROM HOSPITAL DISCHARGE FOR WHO FC II-IV: NO STEP. NEW STARTS FOR WHO FC III-IV: NO STEP. NEW STARTS FOR WHO FC II: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING AGENTS FROM DIFFERENT DRUG CLASSES: 1) FORMULARY VERSION OF AN ORAL ENDOTHELIAL RECEPTOR ANTAGONIST, 2) FORMULARY VERSION OF AN ORAL PHOSPHODIESTERASE TYPE-5 INHIBITOR, OR 3) FORMULARY VERSION OF AN ORAL CGMP STIMULATOR. RENEWAL: IMPROVEMENT FROM BASELINE IN THE 6-MINUTE WALK DISTANCE TEST OR REMAINED STABLE FROM BASELINE IN THE 6-MINUTE WALK DISTANCE TEST AND WHO FC HAS IMPROVED OR REMAINED STABLE. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TRIENTINE

## Products Affected

- *trientine oral capsule 250 mg*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	WILSONS DISEASE: INITIAL: KNOWN FAMILY HISTORY OF WILSONS DISEASE OR PHYSICAL EXAMINATION CONSISTENT WITH WILSONS DISEASE. CONFIRMATION OF ONE OF THE FOLLOWING: 1) PLASMA COPPER-PROTEIN CERULOPLASMIN LESS THAN 20 MG/DL, 2) LIVER BIOPSY POSITIVE FOR AN ABNORMALLY HIGH CONCENTRATION OF COPPER (GREATER THAN 250 MCG/G DRY WEIGHT) OR THE PRESENCE OF KAYSER-FLEISCHER RINGS, OR 3) CONFIRMATION BY GENETIC TESTING FOR ATP7B MUTATIONS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	WILSONS DISEASE: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A HEPATOLOGIST OR GASTROENTEROLOGIST.
<b>Coverage Duration</b>	INITIAL: 12 MONTHS, RENEWAL: LIFETIME.
<b>Other Criteria</b>	WILSONS DISEASE: INITIAL: TRIAL OF OR CONTRAINDICATION TO FORMULARY VERSION OF PENICILLAMINE (DEPEN). RENEWAL: PATIENT CONTINUES TO BENEFIT FROM THE MEDICATION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TRIFLURIDINE/TIPIRACIL

---

## Products Affected

- LONSURF ORAL TABLET 15-6.14 MG, 20-8.19 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# TRIPTORELIN-TRELSTAR

---

## Products Affected

- TRELSTAR INTRAMUSCULAR  
SUSPENSION FOR  
RECONSTITUTION

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS.
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# TRUQAP

## Products Affected

- TRUQAP

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative, locally advanced or metastatic breast cancer with 1 or more PIK3CA/AKT1/PTEN-alterations as detected by an FDA-approved test and, A.) patient has had disease progression following 1 or more endocrine-based regimen(s) in the metastatic setting or recurrence on or within 12 months of completing adjuvant therapy, and B.) will be used in combination with fulvestrant injection.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

# TUCATINIB

---

## Products Affected

- TUKYSA ORAL TABLET 150 MG, 50 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# UBROGEPANT

## Products Affected

- UBRELVY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	ACUTE MIGRAINE TREATMENT: INITIAL: TRIAL OF OR CONTRAINDICATION TO ONE TRIPTAN (E.G., SUMATRIPTAN, RIZATRIPTAN). RENEWAL: 1) IMPROVEMENT FROM BASELINE IN A VALIDATED ACUTE TREATMENT PATIENT-REPORTED OUTCOME QUESTIONNAIRE, OR 2) THERAPY WORKS CONSISTENTLY IN MAJORITY OF MIGRAINE ATTACKS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# UPADACITINIB

## Products Affected

- RINVOQ

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	NON-RADIOGRAPHIC AXIAL SPONDYLOARTHRITIS (NR-AXSPA): 1) C-REACTIVE PROTEIN LEVELS ABOVE THE UPPER LIMIT OF NORMAL, OR 2) SACROILIITIS ON MAGNETIC RESONANCE IMAGING (MRI).
Age Restrictions	
Prescriber Restrictions	INITIAL: RA, AS, NR-AXSPA: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSA: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR DERMATOLOGIST. AD: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST, ALLERGIST, OR IMMUNOLOGIST. UC: PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
Other Criteria	<p>INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) - IF A PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE GREATER THAN OR EQUAL TO 20MG PER WEEK OR MAXIMALLY TOLERATED DOSE IS REQUIRED. PSA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD. ATOPIC DERMATITIS: 1) ATOPIC DERMATITIS COVERING AT LEAST 10 PERCENT OF BODY SURFACE AREA OR ATOPIC DERMATITIS AFFECTING THE FACE, HEAD, NECK, HANDS, FEET, GROIN, OR INTERTRIGINOUS AREAS, 2) INTRACTABLE PRURITUS OR CRACKING/OOZING/BLEEDING OF AFFECTED SKIN, 3) TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING: TOPICAL CORTICOSTEROID, TOPICAL CALCINEURIN INHIBITOR, TOPICAL PDE4 INHIBITOR, OR TOPICAL JAK INHIBITOR, AND 4) NO CONCURRENT USE WITH OTHER SYSTEMIC BIOLOGIC/JAK INHIBITOR FOR THE TREATMENT OF ATOPIC DERMATITIS. UC: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A CORTICOSTEROID (E.G., BUDESONIDE, METHYLPREDNISOLONE), AZATHIOPRINE, MERCAPTOPURINE, METHOTREXATE, OR MESALAMINE. NR-AXSPA: TRIAL OF OR CONTRAINDICATION TO AN NSAID (NON-STEROIDAL ANTI-INFLAMMATORY DRUG). RENEWAL: RA, PSA, AS, NR-AXSPA: CONTINUES TO BENEFIT FROM THE MEDICATION. ATOPIC DERMATITIS: 1) IMPROVEMENT WHILE ON THERAPY, AND 2) NO CONCURRENT USE WITH OTHER SYSTEMIC BIOLOGIC/JAK INHIBITOR FOR THE TREATMENT OF ATOPIC DERMATITIS.</p>
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# USTEKINUMAB

## Products Affected

- STELARA SUBCUTANEOUS

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, GENITAL AREA, OR FACE.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	INITIAL: PSA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG). PSO: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A PUVA (PHOTOTHERAPY ULTRAVIOLET LIGHT A), UVB (ULTRAVIOLET LIGHT B), TOPICAL CORTICOSTEROIDS, CALCIPOTRIENE, ACITRETIN, METHOTREXATE, OR CYCLOSPORINE. CD, UC: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A CORTICOSTEROID (E.G., BUDESONIDE, METHYLPREDNISOLONE), AZATHIOPRINE, MERCAPTOPURINE, METHOTREXATE, OR MESALAMINE. RENEWAL: PSA, PSO: CONTINUES TO BENEFIT FROM THE MEDICATION.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# VALBENAZINE

## Products Affected

- INGREZZA
- INGREZZA INITIATION PACK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	TARDIVE DYSKINESIA (TD): PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST, PSYCHIATRIST, OR MOVEMENT DISORDER SPECIALIST.
Coverage Duration	12 MONTHS
Other Criteria	TD: 1) PRIOR HISTORY OF USING AGENTS THAT CAUSE TARDIVE DYSKINESIA, AND 2) TRIAL OF OR CONTRAINDICATION TO THE PREFERRED AGENT: AUSTEDO.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# VANDETANIB

---

## Products Affected

- CAPRELSA ORAL TABLET 100 MG, 300 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	CURRENTLY STABLE ON CAPRELSA REQUIRES NO EXTRA CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# VANFLYTA

---

## Products Affected

- VANFLYTA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Patient must have all of the following A.) Newly diagnosed acute myeloid leukemia with FLT3-ITD mutation, B.) Used in combination with standard cytarabine and anthracycline induction and cytarabine consolidation, and as maintenance monotherapy following consolidation chemotherapy, and C.) Must be enrolled in the VANFLYTA REMS program
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	12 months
Other Criteria	None
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

# VEMURAFENIB

---

## Products Affected

- ZELBORAF

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	MELANOMA: ZELBORAF WILL BE USED ALONE OR IN COMBINATION WITH COTELLIC
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# VENETOCLAX

---

## Products Affected

- VENCLEXTA ORAL TABLET 10 MG, 100 MG, 50 MG
- VENCLEXTA STARTING PACK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# VERQUVO

## Products Affected

- VERQUVO

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Concomitant use of other soluble guanylate cyclase (sGC) stimulators, or B.) Pregnancy
Required Medical Information	Diagnosis of chronic heart failure (HF), NYHA Class II to IV and all of the following 1.) Left ventricular ejection fraction less than 45%, 2.) Previous hospitalization for HF within 6 months or outpatient IV diuretic treatment for HF within 3 months
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a cardiologist
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

# VIGABATRIN

## Products Affected

- *vigabatrin*
- *vigadrone*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	REFRACTORY COMPLEX PARTIAL SEIZURES (CPS), INFANTILE SPASMS: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	CPS: 1) TRIAL OF OR CONTRAINDICATION TO TWO ANTIEPILEPTIC AGENTS AND 2) BENEFITS OUTWEIGH THE POTENTIAL FOR VISION LOSS. INFANTILE SPASMS: BENEFITS OUTWEIGH THE POTENTIAL FOR VISION LOSS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# VISMODEGIB

---

## Products Affected

- ERIVEDGE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# VORICONAZOLE SUSPENSION

## Products Affected

- *voriconazole oral suspension for reconstitution*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	CANDIDA INFECTIONS: 3 MOS. ALL OTHER INDICATIONS: 6 MOS.
Other Criteria	CANDIDA INFECTIONS: TRIAL OF OR CONTRAINDICATION TO FLUCONAZOLE. ALL INDICATIONS: INABILITY TO SWALLOW TABLETS OR AN INDICATION FOR ESOPHAGEAL CANDIDIASIS. CONTINUATION OF THERAPY AFTER HOSPITAL DISCHARGE REQUIRES NO EXTRA CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# ZANUBRUTINIB

---

## Products Affected

- BRUKINSA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# ZURZUVAE

---

## Products Affected

- ZURZUVAE ORAL CAPSULE 20 MG, 25 MG, 30 MG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of postpartum depression
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	14 days
Other Criteria	None
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

## INDEX

<i>abiraterone</i> .....	4	BENLYSTA INTRAVENOUS.....	31
ACTEMRA.....	301, 303	BENLYSTA SUBCUTANEOUS.....	31
ACTEMRA ACTPEN.....	303	BESREMI.....	255
ACTHAR.....	59	<i>betaine</i> .....	37
ACTIMMUNE.....	153	BETASERON SUBCUTANEOUS KIT	
ADAKVEO.....	61	.....	151
<i>adalimumab-fkjp</i> .....	8	<i>bexarotene</i> .....	38
ADEMPAS.....	247	<i>bortezomib injection recon soln 1 mg, 2.5</i>	
AJOVY AUTOINJECTOR.....	124	<i>mg</i> .....	40
AJOVY SYRINGE.....	124	<i>bosentan</i> .....	41
ALECENSA.....	12	BOSULIF ORAL TABLET 100 MG,	
ALTRENO.....	308	400 MG, 500 MG.....	43
ALUNBRIG ORAL TABLET 180 MG,		BRAFTOVI.....	100
30 MG, 90 MG.....	44	BRONCHITOL.....	45
ALUNBRIG ORAL TABLETS,DOSE		BRUKINSA.....	333
PACK.....	44	CABLIVI INJECTION KIT.....	50
<i>alyq</i> .....	218	CABOMETYX ORAL TABLET 20	
<i>ambrisentan</i> .....	14	MG, 40 MG, 60 MG.....	48
AMJEVITA(CF) SUBCUTANEOUS		CALQUENCE.....	6
SYRINGE 10 MG/0.2 ML.....	8	CALQUENCE (ACALABRUTINIB	
<i>apomorphine</i> .....	17	MAL).....	6
<i>armodafinil</i> .....	194	CAPRELSA ORAL TABLET 100 MG,	
AUBAGIO.....	293	300 MG.....	325
AUSTEDO ORAL TABLET 12 MG, 6		<i>carglumic acid</i> .....	52
MG, 9 MG.....	79	CAYSTON.....	27
AUSTEDO XR ORAL TABLET		CERDELGA.....	97
EXTENDED RELEASE 24 HR 12		CIMZIA.....	54
MG, 24 MG, 6 MG.....	79	CIMZIA POWDER FOR RECONST...54	
AUSTEDO XR TITRATION		COMETRIQ ORAL CAPSULE 100	
KT(WK1-4).....	79	MG/DAY(80 MG X1-20 MG X1), 140	
<i>avita topical gel</i> .....	308	MG/DAY(80 MG X1-20 MG X3), 60	
AVONEX INTRAMUSCULAR PEN		MG/DAY (20 MG X 3/DAY).....	47
INJECTOR KIT.....	150	COPAXONE SUBCUTANEOUS	
AVONEX INTRAMUSCULAR		SYRINGE 20 MG/ML, 40 MG/ML....	132
SYRINGE KIT.....	150	COPIKTRA.....	89
AVSOLA.....	148	CORTROPHIN GEL.....	59
AYVAKIT.....	23	COSENTYX (2 SYRINGES).....	262
BALVERSA ORAL TABLET 3 MG, 4		COSENTYX INTRAVENOUS.....	262
MG, 5 MG.....	106	COSENTYX PEN (2 PENS).....	262
<i>bendamustine intravenous recon soln</i> .....	34	COSENTYX SUBCUTANEOUS	
BENDAMUSTINE INTRAVENOUS		SYRINGE 75 MG/0.5 ML.....	262
SOLUTION.....	34	COSENTYX UNOREADY PEN.....	262
BENDEKA.....	34	COTELLIC.....	58



CYLTEZO(CF).....	8	ENSPRYNG.....	261
CYLTEZO(CF) PEN.....	8	ENTADFI.....	118
CYLTEZO(CF) PEN CROHN'S-UC- HS.....	8	EPCLUSA ORAL PELLETS IN PACKET 150-37.5 MG, 200-50 MG.....	272
CYLTEZO(CF) PEN PSORIASIS-UV....	8	EPCLUSA ORAL TABLET.....	272
CYSTADROPS.....	63	EPIDIOLEX.....	49
CYSTARAN.....	63	<i>epoprostenol</i> .....	105
<i>dalfampridine</i> .....	66	ERIVEDGE.....	331
DAURISMO ORAL TABLET 100 MG, 25 MG.....	131	ERLEADA ORAL TABLET 240 MG, 60 MG.....	16
DAYBUE.....	70	<i>erlotinib oral tablet 100 mg, 150 mg, 25 mg</i> .....	107
<i>deferasirox</i> .....	72	EVENITY SUBCUTANEOUS SYRINGE 210MG/2.34ML (	
<i>deferiprone</i> .....	74	105MG/1.17MLX2).....	253
<i>deferoxamine injection recon soln 500 mg</i> ..	76	<i>everolimus (antineoplastic) oral tablet 10 mg, 2.5 mg, 5 mg, 7.5 mg</i> .....	111
DIACOMIT ORAL CAPSULE 250 MG, 500 MG.....	283	<i>everolimus (antineoplastic) oral tablet for suspension</i> .....	111
DIACOMIT ORAL POWDER IN PACKET 250 MG, 500 MG.....	283	EVRYSDI.....	252
<i>diclofenac sodium topical gel 3 %</i> .....	80	EXKIVITY.....	191
<i>diclofenac sodium topical solution in metered-dose pump</i> .....	81	FARYDAK.....	214
<i>dimethyl fumarate oral capsule, delayed release(drlec) 120 mg, 120 mg (14)- 240 mg (46), 240 mg</i> .....	82	FASENRA.....	35
DOPTELET (10 TAB PACK).....	24	FASENRA PEN.....	35
DOPTELET (15 TAB PACK).....	24	<i>fentanyl citrate buccal lozenge on a handle</i> .....	114
DOPTELET (30 TAB PACK).....	24	FERRIPROX (2 TIMES A DAY).....	74
<i>dronabinol</i> .....	84	FERRIPROX ORAL SOLUTION.....	74
<i>droxidopa</i> .....	85	<i> fingolimod</i> .....	120
DUPIXENT PEN.....	86	FINTEPLA.....	113
DUPIXENT SYRINGE.....	86	FIRMAGON KIT W DILUENT SYRINGE SUBCUTANEOUS	
EGRIFTA SV.....	294	RECON SOLN 120 MG, 80 MG.....	122
ELIGARD.....	171	FOTIVDA.....	300
ELIGARD (3 MONTH).....	171	FRUZAQLA ORAL CAPSULE 1 MG, 5 MG.....	125
ELIGARD (4 MONTH).....	171	FULPHILA.....	222
ELIGARD (6 MONTH).....	171	FYLNETRA.....	219
EMGALITY PEN.....	127	GALAFOLD.....	188
EMGALITY SYRINGE SUBCUTANEOUS SYRINGE 120 MG/ML, 300 MG/3 ML (100 MG/ML X 3).....	127	GATTEX 30-VIAL.....	290
ENBREL.....	109	GAVRETO.....	237
ENBREL MINI.....	109	<i>gefitinib</i> .....	129
ENBREL SURECLICK.....	109	GILENYA ORAL CAPSULE 0.25 MG .....	120
ENDARI.....	175	GILOTRIF.....	11

<i>glatiramer subcutaneous syringe 20 mg/ml, 40 mg/ml</i> .....	132	<i>ibuprofen-famotidine</i> .....	142
<i>glatopa subcutaneous syringe 20 mg/ml, 40 mg/ml</i> .....	132	<i>icatibant</i> .....	143
GRANIX.....	289	ICLUSIG.....	234
HADLIMA.....	8	IDHIFA.....	99
HADLIMA PUSHTOUCH.....	8	ILUMYA.....	299
HADLIMA(CF).....	8	<i>imatinib oral tablet 100 mg, 400 mg</i> .....	145
HADLIMA(CF) PUSHTOUCH.....	8	IMBRUVICA ORAL CAPSULE 140 MG, 70 MG.....	141
HAEGARDA SUBCUTANEOUS RECON SOLN 2,000 UNIT, 3,000 UNIT.....	46	IMBRUVICA ORAL SUSPENSION... 141	
HARVONI ORAL PELLETS IN PACKET 33.75-150 MG, 45-200 MG... 165		IMBRUVICA ORAL TABLET.....	141
HARVONI ORAL TABLET.....	165	IMPAVIDO.....	190
HETLIOZ.....	287	INBRIJA INHALATION CAPSULE, W/INHALATION DEVICE.....	174
HETLIOZ LQ.....	287	<i>infliximab</i> .....	146
HULIO(CF).....	8	INGREZZA.....	324
HULIO(CF) PEN.....	8	INGREZZA INITIATION PACK.....	324
HUMIRA PEN.....	8	INLYTA ORAL TABLET 1 MG, 5 MG.....	25
HUMIRA PEN CROHNS-UC-HS STARTER.....	8	INQOVI.....	71
HUMIRA PEN PSOR-UEITS-ADOL HS.....	8	INREBIC.....	112
HUMIRA SUBCUTANEOUS SYRINGE KIT 40 MG/0.8 ML.....	8	IRESSA.....	129
HUMIRA(CF).....	8	<i>itraconazole oral solution</i> .....	154
HUMIRA(CF) PEDI CROHNS STARTER.....	8	<i>ivermectin oral</i> .....	156
HUMIRA(CF) PEN.....	8	JAKAFI.....	257
HUMIRA(CF) PEN CROHNS-UC-HS... 8		<i>javygtor oral tablet, soluble</i> .....	259
HUMIRA(CF) PEN PEDIATRIC UC... 8		JAYPIRCA ORAL TABLET 100 MG, 50 MG.....	232
HUMIRA(CF) PEN PSOR-UV-ADOL HS.....	8	JUXTAPID ORAL CAPSULE 10 MG, 20 MG, 30 MG, 5 MG.....	178
HYRIMOZ PEN CROHN'S-UC STARTER.....	8	JYNARQUE ORAL TABLET.....	307
HYRIMOZ PEN PSORIASIS STARTER.....	8	JYNARQUE ORAL TABLETS, SEQUENTIAL.....	307
HYRIMOZ(CF).....	8	KALYDECO ORAL GRANULES IN PACKET 25 MG, 5.8 MG, 50 MG, 75 MG.....	155
HYRIMOZ(CF) PEDI CROHN STARTER SUBCUTANEOUS SYRINGE 80 MG/0.8 ML- 40 MG/0.4 ML.....	8	KALYDECO ORAL TABLET.....	155
HYRIMOZ(CF) PEN.....	8	KERENDIA.....	119
IBRANCE.....	213	KESIMPTA PEN.....	201
		KEVZARA.....	260
		KINERET.....	15
		KISQALI FEMARA CO-PACK ORAL TABLET 200 MG/DAY(200 MG X 1)- 2.5 MG, 400 MG/DAY(200 MG X 2)- 2.5 MG, 600 MG/DAY(200 MG X 3)- 2.5 MG.....	243

KISQALI ORAL TABLET 200 MG/DAY (200 MG X 1), 400 MG/DAY (200 MG X 2), 600 MG/DAY (200 MG X 3).....	243	MAYZENT ORAL TABLET 0.25 MG, 1 MG, 2 MG.....	269
KORLYM.....	187	MAYZENT STARTER(FOR 1MG MAINT).....	269
KOSELUGO ORAL CAPSULE 10 MG, 25 MG.....	268	MAYZENT STARTER(FOR 2MG MAINT).....	269
KRAZATI.....	7	MEKINIST ORAL TABLET 0.5 MG, 2 MG.....	309
KYNMOBI SUBLINGUAL FILM 10 MG, 10-15-20-25-30 MG, 15 MG, 20 MG, 25 MG, 30 MG.....	18	MEKTOVI.....	39
<i>lanreotide</i> .....	162	<i>miglustat</i> .....	189
<i>lapatinib</i> .....	163	<i>modafinil oral tablet 100 mg, 200 mg</i> .....	194
<i>lenalidomide</i> .....	166	<i>morphine concentrate oral solution</i> .....	139
LENVIMA.....	167	NATPARA.....	215
<i>leuprolide (3 month)</i> .....	170	NERLYNX.....	195
<i>leuprolide subcutaneous kit</i> .....	169	NEULASTA.....	219
<i>lidocaine hcl mucous membrane solution 4</i> <i>% (40 mg/ml)</i> .....	176	NINLARO.....	158
<i>lidocaine topical adhesive patch,medicated</i> <i>5 %</i> .....	176	<i>nitisinone oral capsule 10 mg, 2 mg, 5 mg</i> .....	199
<i>lidocaine topical ointment</i> .....	176	NIVESTYM.....	115
<i>lidocaine-prilocaine topical cream</i> .....	177	NORDITROPIN FLEXPPO.....	276
LONSURF ORAL TABLET 15-6.14 MG, 20-8.19 MG.....	315	NOXAFIL INTRAVENOUS.....	236
LORBRENA ORAL TABLET 100 MG, 25 MG.....	180	NOXAFIL ORAL SUSP,DELAYED RELEASE FOR RECON.....	236
LUMAKRAS.....	282	NOXAFIL ORAL SUSPENSION.....	235
LUPRON DEPOT.....	172	NUBEQA.....	67
LUPRON DEPOT (3 MONTH).....	172	NUCALA SUBCUTANEOUS AUTO- INJECTOR.....	183
LUPRON DEPOT (4 MONTH).....	172	NUCALA SUBCUTANEOUS RECON SOLN.....	183
LUPRON DEPOT (6 MONTH).....	172	NUCALA SUBCUTANEOUS SYRINGE 100 MG/ML, 40 MG/0.4 ML.....	183
LUPRON DEPOT-PED INTRAMUSCULAR KIT 11.25 MG...	172	NUPLAZID.....	230
LYBALVI.....	203	NURTEC ODT.....	245
LYNPARZA.....	204	NYVEPRIA.....	220
LYTGOBI.....	126	OCALIVA.....	200
MAVENCLAD (10 TABLET PACK)....	56	ODOMZO.....	280
MAVENCLAD (4 TABLET PACK).....	56	OFEV.....	197
MAVENCLAD (5 TABLET PACK).....	56	OJJAARA.....	202
MAVENCLAD (6 TABLET PACK).....	56	OLUMIANT.....	28
MAVENCLAD (7 TABLET PACK).....	56	ONGENTYS.....	210
MAVENCLAD (8 TABLET PACK).....	56	ONUREG.....	26
MAVENCLAD (9 TABLET PACK).....	56	OPSUMIT.....	182
MAVYRET.....	133	ORENCIA.....	1
		ORENCIA CLICKJECT.....	1
		ORFADIN ORAL CAPSULE 20 MG.	199

ORFADIN ORAL SUSPENSION.....	199	REGRANEX.....	29
ORGOVYX.....	242	RELEUKO SUBCUTANEOUS.....	116
ORILISSA ORAL TABLET 150 MG, 200 MG.....	93	RELISTOR ORAL.....	185
ORKAMBI ORAL GRANULES IN PACKET.....	181	RELISTOR SUBCUTANEOUS SOLUTION.....	184
ORKAMBI ORAL TABLET.....	181	RELISTOR SUBCUTANEOUS SYRINGE 12 MG/0.6 ML, 8 MG/0.4 ML.....	184
ORSERDU ORAL TABLET 345 MG, 86 MG.....	92	RETACRIT INJECTION SOLUTION 10,000 UNIT/ML, 2,000 UNIT/ML, 20,000 UNIT/2 ML, 20,000 UNIT/ML, 3,000 UNIT/ML, 4,000 UNIT/ML, 40,000 UNIT/ML.....	103
OTEZLA.....	19	RETEVMO ORAL CAPSULE 40 MG, 80 MG.....	267
OTEZLA STARTER.....	19	REVCovi.....	95
<i>oxycodone oral concentrate</i> .....	139	REZLIDHIA.....	205
OZEMPIC SUBCUTANEOUS PEN INJECTOR 1 MG/DOSE (4 MG/3 ML), 2 MG/DOSE (8 MG/3 ML).....	135	REZUROCK.....	32
PALYNZIQ.....	224	RINVOQ.....	320
<i>pazopanib</i> .....	217	ROZLYTREK ORAL CAPSULE 100 MG, 200 MG.....	101
PEMAZYRE.....	226	ROZLYTREK ORAL PELLETS IN PACKET.....	101
<i>penicillamine oral tablet</i> .....	227	RUBRACA.....	256
PIQRAY ORAL TABLET 200 MG/DAY (200 MG X 1), 250 MG/DAY (200 MG X1-50 MG X1), 300 MG/DAY (150 MG X 2).....	13	RYBELSUS.....	135
<i>pirfenidone oral capsule</i> .....	231	RYDAPT.....	186
<i>pirfenidone oral tablet 267 mg, 534 mg, 801 mg</i> .....	231	SAIZEN SUBCUTANEOUS RECON SOLN 5 MG.....	278
PLEGRIDY SUBCUTANEOUS PEN INJECTOR 125 MCG/0.5 ML, 63 MCG/0.5 ML- 94 MCG/0.5 ML.....	152	<i>sajazir</i> .....	143
PLEGRIDY SUBCUTANEOUS SYRINGE 125 MCG/0.5 ML, 63 MCG/0.5 ML- 94 MCG/0.5 ML.....	152	<i>sapropterin oral tablet, soluble</i> .....	259
POMALYST.....	233	SCEMBLIX.....	21
<i>posaconazole oral</i> .....	235	SEROSTIM SUBCUTANEOUS RECON SOLN 4 MG, 5 MG, 6 MG....	278
PREVYMIS ORAL.....	168	SIGNIFOR.....	216
PROMACTA ORAL POWDER IN PACKET 12.5 MG, 25 MG.....	98	SIKLOS ORAL TABLET 100 MG.....	140
PROMACTA ORAL TABLET 12.5 MG, 25 MG, 50 MG, 75 MG.....	98	<i>sildenafil (pulm.hypertension) intravenous</i> .....	218
<i>pyrimethamine</i> .....	239	<i>sildenafil (pulm.hypertension) oral tablet</i> .....	218
QINLOCK.....	249	SIRTURO.....	30
<i>quinine sulfate</i> .....	240	SKYRIZI INTRAVENOUS.....	250
QULIPTA.....	22	SKYRIZI SUBCUTANEOUS PEN INJECTOR.....	250
RADICAVA.....	90	SKYRIZI SUBCUTANEOUS SYRINGE 150 MG/ML.....	250
RAVICTI.....	136		

SKYRIZI SUBCUTANEOUS		TIBSOVO.....	157
WEARABLE INJECTOR.....	250	TRACLEER ORAL TABLET FOR	
<i>sodium oxybate</i> .....	270	SUSPENSION.....	41
SOMATULINE DEPOT		TRELSTAR INTRAMUSCULAR	
SUBCUTANEOUS SYRINGE 120		SUSPENSION FOR	
MG/0.5 ML.....	162	RECONSTITUTION.....	316
SOMAVERT.....	225	TREMFYA.....	138
<i>sorafenib</i> .....	281	<i>treprostinil sodium</i> .....	312
SPRAVATO.....	108	<i>tretinoin</i> .....	308
SPRYCEL ORAL TABLET 100 MG,		<i>trientine oral capsule 250 mg</i> .....	314
140 MG, 20 MG, 50 MG, 70 MG, 80		TRIKAFTA ORAL TABLETS,	
MG.....	69	SEQUENTIAL.....	96
STELARA SUBCUTANEOUS.....	322	TRULICITY.....	135
STIVARGA.....	241	TRUQAP.....	317
<i>sunitinib malate</i> .....	284	TUKYSA ORAL TABLET 150 MG, 50	
SUNOSI.....	275	MG.....	318
SYMDEKO.....	297	TURALIO ORAL CAPSULE 200 MG	229
SYMLINPEN 120.....	238	TYVASO.....	310
SYMLINPEN 60.....	238	UBRELVY.....	319
SYMPAZAN.....	57	UDENYCA.....	221
SYNAREL.....	192	UDENYCA AUTOINJECTOR.....	221
SYNRIBO.....	206	UPTRAVI ORAL TABLET 1,000	
TABRECTA.....	51	MCG, 1,200 MCG, 1,400 MCG, 1,600	
<i>tadalafil (pulm. hypertension)</i> .....	218	MCG, 200 MCG, 400 MCG, 600 MCG,	
<i>tadalafil oral tablet 2.5 mg, 5 mg</i> .....	285	800 MCG.....	264
TAFINLAR ORAL CAPSULE.....	64	UPTRAVI ORAL TABLETS,DOSE	
TAGRISSO.....	211	PACK.....	264
TAKHZYRO SUBCUTANEOUS		VANFLYTA.....	326
SOLUTION.....	161	VENCLEXTA ORAL TABLET 10	
TAKHZYRO SUBCUTANEOUS		MG, 100 MG, 50 MG.....	328
SYRINGE 150 MG/ML, 300 MG/2 ML		VENCLEXTA STARTING PACK.....	328
(150 MG/ML).....	161	VERQUVO.....	329
TALTZ AUTOINJECTOR.....	159	VERZENIO.....	3
TALTZ SYRINGE.....	159	<i>vigabatrin</i> .....	330
TALZENNA.....	286	<i>vigadrone</i> .....	330
TASCENSO ODT.....	121	VITRAKVI ORAL CAPSULE 100	
TASIGNA ORAL CAPSULE 150 MG,		MG, 25 MG.....	164
200 MG, 50 MG.....	196	VITRAKVI ORAL SOLUTION.....	164
<i>tasimelteon</i> .....	287	VIZIMPRO.....	65
TAVALISSE.....	123	VONJO.....	212
TAZVERIK.....	288	<i>voriconazole oral suspension for</i>	
TEPMETKO.....	292	<i>reconstitution</i> .....	332
<i>teriflunomide</i> .....	293	VOSEVI.....	273
<i>tetrabenazine</i> .....	296	VOTRIENT.....	217
THALOMID.....	298	VUMERITY.....	83

WELIREG.....	33
XADAGO.....	258
XALKORI ORAL CAPSULE.....	62
XELJANZ.....	305
XELJANZ XR.....	305
XERMELO.....	291
XGEVA.....	78
XIFAXAN ORAL TABLET 200 MG, 550 MG.....	244
XOLAIR SUBCUTANEOUS RECON SOLN.....	207
XOLAIR SUBCUTANEOUS SYRINGE 150 MG/ML, 75 MG/0.5 ML.....	207
XOSPATA.....	130
XPOVIO ORAL TABLET 100 MG/WEEK (50 MG X 2), 40 MG/WEEK (40 MG X 1), 40MG TWICE WEEK (40 MG X 2), 60 MG/WEEK (60 MG X 1), 60MG TWICE WEEK (120 MG/WEEK), 80 MG/WEEK (40 MG X 2), 80MG TWICE WEEK (160 MG/WEEK).....	266
XTANDI ORAL CAPSULE.....	102
XTANDI ORAL TABLET 40 MG, 80 MG.....	102
XYOSTED.....	295
XYREM.....	270
<i>yargesa</i> .....	189
YONSA.....	5
YUSIMRY(CF) PEN.....	8
ZARXIO.....	117
ZEJULA ORAL CAPSULE.....	198
ZEJULA ORAL TABLET.....	198
ZELBORAF.....	327
ZIEXTENZO.....	223
ZOLADEX.....	137
ZOMACTON SUBCUTANEOUS RECON SOLN 5 MG.....	278
ZTALMY.....	128
ZTLIDO.....	176
ZURZUVAE ORAL CAPSULE 20 MG, 25 MG, 30 MG.....	334
ZYDELIG.....	144
ZYKADIA.....	53

